

Evaluation and impact of fetal physiology training on fetal heart rate analysis.

Harmonie Dupuis¹, Louise Ghesquiere², Adeline Pierache³, Damien Subtil⁴, Veronique Houfflin-Debarge³, and Charles Garabedian⁴

¹Jeanne de Flandre Hospital

²CHU Lille Jeanne de Flandre

³CHRU de Lille

⁴Jeanne De Flandre

May 5, 2020

Abstract

Objective - This study aims to assess fetal physiology training in terms of theoretical knowledge, fetal heart rate interpretation and use of second-line examination. **Design** - Single-center prospective study (CHU Lille, France) **Setting** - The evaluation of fetal well-being during labor is based on fetal heart rate (FHR) analysis and requires knowledge of physiology. **Population** - Obstetrics and gynecology residents from November 2017 to November 2018 (n=34) **Methods** - The training was conducted in 3 steps: a session of FHR interpretation and the use of fetal scalp blood sampling (FBS) on clinical cases, then a teaching session on fetal physiology, and finally another session on the same cases presented in the first one. **Main Outcome Measures** - Theoretical knowledge evaluation (MCQs), number of FBS requested, the reproducibility of responses. **Results** - Almost 3% estimated their training sufficient on fetal physiology, 11.8% on fetal heart rate analysis and 14.7% on second-line examination. The training allowed a significant improvement of their theoretical knowledge evaluation (median [IQR] : 1.5[1.0 to 2.0] vs 4.0[3.0 to 4.5] of MCQs, $p < 0.001$), a decrease in the number of FBS requested (36.3% vs 29.5%, $p = 0.002$). The Krippendorff's alpha index assessing the reproducibility of their response was significantly improved, reflecting a better homogenization of practices (alpha [IC95] : 0.60[0.55 to 0.65] vs 0.72[0.67 to 0.76]). **Conclusions** - The improvement of knowledge in fetal physiology allows a better interpretation of the FHR with better indications of second-line examinations and a homogenization of practices. **Funding** - None

Manuscript

Introduction

Fetal surveillance during labor is primarily based on fetal heart rate (FHR) analysis. The purpose of the interpretation of the FHR is to detect signs of poor fetal tolerance and therefore situations at risk of fetal acidosis. Indeed, this acidosis can be responsible for morbidity and perinatal mortality including motor disability of cerebral origin (1). In these intermediate-risk situations of fetal acidosis, second-line methods exist such as fetal scalp sampling (pH or lactate measurement) or ST segment analysis (2–4).

Interpretation of abnormalities observed on the FHR or second-line exams requires knowledge of fetal physiology during labor (5,6). The recent FIGO recommendations modified in 2015 classify the FHR as normal, suspicious or pathological, leading to a more physiological analysis of the FHR. (7). On the other hand, the place of second-line exams is controversial. In fact, a US randomized study concluded that ST segment analysis was not helpful in preventing fetal acidosis and nor in reducing intervention (cesarean or instrumental extraction) (8). Moreover, the physiopathology and the interest of fetal blood scalp sampling are discussed (9). Therefore, it is proposed to improve the quality of the interpretation of the FHR thanks to a

better knowledge of the fetal physiology and of the adaptation of the fetus to the hypoxemia during labor. Thus, the aim of the present study was to assess fetal physiology training in terms of theoretical knowledge, interpretation of FHR and use of second-line examination.

Methods

A single-center prospective study (CHU Lille, France) was conducted among obstetrics and gynecology residents from November 2017 to November 2018 whatever their degree of competence (residency lasts 5 years in France).

The evaluation of the residents was conducted in 3 stages. The first evaluation session was organized before our teaching session called "pre-training period". After this first evaluation session, the residents received one hour and half of teaching on fetal physiology and FHR analysis. The second evaluation session, called "post-training period", was organized after this teaching and these three sessions were realized at 3 different times during their internship.

Each assessment session (pre- and post-training period) was organized into 2 parts: a theoretical part of multiple-choice questions (MCQs), and a practical part on clinical cases. During the theoretical evaluation, the residents had to answer a series of 7 multiple-choice questions. For each question, five answers were proposed and several answers were possible. They got one point to the question if all the answers were correct, 0.5 point if they made a mistake and no point beyond an error. These questions focused on the fetal response to hypoxemia and thus to fetal physiology during labor.

The practical part was based on 6 clinical cases resulting from real clinical situations chosen by 2 investigators of the study. These were cases of singleton pregnancy in labor (spontaneous or induced) at term and with presence of abnormalities of the FHR for which the medical team had been solicited. Among the 6 files selected, 5 had been the subject of one or more FBS for a total of 10 fetal pH measurements actually achieved. For each of these clinical cases, 3 to 5 periods (27 in total) were chosen during the labor and at each period the residents had to analyze the FHR according to the CNGOF classification (10) divided into 5 categories: normal, low risk of acidosis, intermediate risk of acidosis, significant risk of acidosis, major risk. They were then asked if they achieved FBS or not, and the estimate of the result of this pH measurement : below 7.20, between 7.20 and 7.25 or above 7.25 (3). The second evaluation session (after teaching) was based on the same clinical cases, the neonatal outcome was not given at the end of the first session so as not to influence the answers (11).

The teaching session consisted of a 1.5 hour course. During this session the basics of fetal physiology and FHR analysis were recalled. This class was based on the literature on this topic (12,13) and carried out in a team working in experimental development of a new fetal monitoring tool based on the analysis of the autonomic nervous system (14,15).

Statistical analyzes

Qualitative variables were described in terms of frequency and percentage. The note of theoretical radiographic knowledge of the residents has been described in terms of median and interquartile range.

The comparison of the radiographic theoretical knowledge scores of the residents between the two sessions was evaluated using a rank test signed of Wilcoxon.

The distribution of FHR evaluation grades was compared between the 2 sessions using a mixed ordinal regression model including the session as a fixed effect and an internal random effect to take into account the correlation between the different assessments by resident (27 scenarios, 2 sessions). To evaluate the impact of the training on the analyze of the FHR by the residents, the reproducibility of the responses between the 34 residents on the FHR graduation was evaluated using the Krippendorff's alpha coefficient with its confidence interval of 95% [95% CI].

The FBS requested rate and the rate of correct estimation of the pH (if requested) were compared between the 2 sessions using a mixed logistic model including the session as a fixed effect and an effect internal

random.

The p-values were calculated with a significance level of 5%. Statistical analyzes were performed using the SAS software (SAS Institute version 9.4).

Results

A total of 34 residents participated in the 3 sessions, with seniority ranging from the 1st to the 5th year (Table 1). Only 11.8% of the residents surveyed felt that they had sufficient training on FHR analysis and 2.9% on fetal physiology during labor. Of these, 14.7% felt that they had sufficient training in the indication of second-line surveillance examinations. They were 81.8% have received specific training on FHR during their residency, mainly during a planned course in their university studies. Yet 90.9% believed that their training was insufficient on the use of the FIGO and CNGOF classifications. As for FBS, 66.7% had already laid an indication and 48.4% had never achieved a FBS.

The median MCQ score before teaching was 1.5 [1.0 to 2.0] with scores ranging from 0.0 to 4.5. After our training, the median was 4.0 [3.0 to 4.5] with scores ranging from 1.0 to 6.5 ($p < 0.0001$). Of all the FHR analyzed during the defined periods, 49.2% of the rhythms were considered normal or low risk before our training against 54.7% after (Table 2). The distribution of FHR grades found a less pejorative evaluation after training ($p < 0.001$). In fact, 17.4% of the FHR analyzed by the residents were considered to be at significant risk of acidosis and 2.7% at major risk in pre-training, compared with respectively 11.2% and 0.1%. The reproducibility of residents' responses to FHR evaluation was improved after training with Krippendorff's alpha index pre-training of 0.60 [IC95, 0.55 to 0.65] and post training of 0.72 [IC 9, 0.67 to 0.76]. Before our formation, 327 FBS were requested on all clinical cases against 269 after our training, a decrease of 6.9% ($p = 0.002$). The evaluation of the pH estimate relative to the actual value was significantly different between the two sessions ($p = 0.020$) with a higher good estimate rate after training (47.2% vs 40.0%) although not significant with $p = 0.28$, a lower underestimation rate after training (26.4% vs. 38.2%) and a higher overestimate rate after training (26.4% vs. 21.8%). The rate of underestimation and overestimation was identical after training.

Discussion

Main Findings

The cardiotocograph was developed in the 1960s to improve fetal surveillance. But its generalization has led to an increase in the rate of cesarean section and extractions for FHR abnormalities and without significant reduction in neonatal risk (16,17). A recent Cochrane review also found no differences in cerebral palsy, infant mortality, or other standard measures of neonatal well-being between intermittent or continuous FHR auscultation (17). The only difference found is the reduction of neonatal convulsion rates during continuous auscultation. These findings, including the increase in caesarean section rate, are related to the high sensitivity of the FHR and its low specificity, but not only. Misinterpretations or erroneous decisions are involved with a failure to take into account a pathological pattern in 20% of cases in newborns with metabolic acidosis (18). It therefore appears essential to improve our analysis of FHR and we wanted to evaluate the interest of training on fetal physiology in obstetric gynecology residents regardless of their initial level. We find a better global knowledge in fetal physiology with a reduction of "extreme" classification during the analysis of FHR, resulting in a decrease in the number of FBS, a better estimation of these and a homogenization of practices in clinical cases performed.

The FHR interpretation training starts at the residency but the training time is variable and generally limited. The training is not systematic in France and varies according to the enrollments in workshops during congresses and the courses planned in the university. Conversely, in other countries such as England, the regulation has imposed the need for continuing education on the interpretation of the FHR every 6 months for midwives (19). Therefore, the formation of residents on the FHR and fetal physiology is insufficient in France and all of our residents interviewed were applicants for such training.

The FHR analysis makes it possible to detect situations at risk of fetal acidosis when its interpretation is correctly performed and thus to provide indications of second-line examinations or fetal extraction. Metabolic acidosis and associated neonatal morbidity could potentially be prevented in 40-50% of cases (18). Indeed, the most found errors are a misinterpretation of the RCF, an imprudent use of oxytocin and a failure to recognize at-risk pregnancies (20–22). Training development could reduce the consequences of inadequate monitoring. Indeed, Draycott and al conducted a retrospective study evaluating the value of training in obstetric emergencies. They were interested in the Apgar scores at 5 minutes of all the liveborn singletons with vaginal deliveries at term, between 1998 and 2003. They also identified hypoxic and ischemic encephalopathies. All the medical staff (midwife, gynecologist, anesthesiologist ...) benefited from a training day during the year 2000. Draycott and al did not analyze the year 2000 and compared two periods: a pre-training period (from 1998 to 1999) and a post-training period (from 2001 to 2003). They found a significant reduction in low Apgar scores (<6) and in the incidence of hypoxic and ischemic encephalopathy (23). Thellesen and al, for their part, found a 14% decrease in fetal extraction, without increased risk of fetal hypoxia, after a training program for midwives and gynecologists from a Danish maternity hospital (24). Their training consisted of e-learning sessions and a day of theoretical courses.

The interpretation of the FHR is subject to intra- and inter-observer variability well studied now (25–27). This variability persists despite the existence of classification and this is more important when it comes to FHR classified as intermediate or pathological according to the FIGO classification (28). Regular training on FHR could reduce this inter-observer variability. Pehrson and al interviewed the Medline database to study and evaluate FHR training programs (29). Of the 409 citations they found, 20 studies included and analyzed. They report a better inter-observer agreement after training on the physiology and interpretation of FHR. We had the same findings by studying the reproducibility of the responses and therefore the inter-observer variability via the Krippendorff's alpha index (0.60 before training vs. 0.72 after). Therefore, training in fetal physiology promotes the homogenization of answers. Thus, an education in the interpretation of the FHR would allow a reduction of its variability inter and intra observer and therefore standardize our practices.

Finally, we wanted to evaluate the impact on the use of a second-line examination. Indeed, when the FHR analysis is considered non-reassuring, there are different second-line exams to better characterize the fetal state such as the FBS with pH measurement to study the acidobasic state of the fetus (3,4,30). The interest of FBS is currently debated. It may not be representative of fetal acid-base status as it is derived from peripheral tissue or because of the compression of the fetal scalp during labor for example. This could lead to unnecessary interventions in fetuses that are not really hypoxic, which means that the debate on this subject persists (9,31). The purpose of this study was not to discuss the interest of FBS but to evaluate the decision to perform this second-line exam in our residents. We note a decrease in the number of FBS requested so second-line examination, currently being discussed, through a better interpretation.

Strengths and Limitations

Our study was conducted directly with our residents and the clinical cases were derived from real situations in order to get as close as possible to a current clinical practice. However neonatal outcome was not communicated at the end of the first session so as not to influence the responses of the residents, the knowledge of an unfavorable neonatal outcome leading to a more pessimistic evaluation of the FHR (11,32).

This study has limitations. Only 6 cases were selected reflecting only part of FHR abnormalities and obstetric context. In addition, during the second session, the same clinical cases were evaluated. Finally, these positive results are based on theoretical cases and it will be interesting to evaluate their impact in current practice. We also choose not to ask experts to classify fetal heart rate. Even if interpretation of FHR was more homogenous after the formation, we cannot conclude on a better practice.

Interpretation

In this study, we showed that it is necessary to improve our knowledge of fetal physiology for a better interpretation of FHR with better indications of FBS and more generally, more similar practices between practitioners.

Conclusion

Improved knowledge in fetal physiology allows for better interpretation of FHR with better second-line examination indications and homogenization of practices. It will be interesting to evaluate the impact of training of all professionals in current practice.

Word count

Manuscript: 2298

Introduction: 238

Discussion, conclusion: 1051

Acknowledgements

Disclosure of Interests

The authors report no conflict of interest.

Contribution to Authorship

HD, LG and CG performed the evaluation of the residents and wrote up. CG, DS, and VHD contributed to the conception of this study and to the organization of this one with the residents. Statistical analyzes were carried out by AP.

Details of Ethics Approval

Declaration N/Réf : CNIL DEC19-500, January 27th 2020

Funding

None

References

1. Boog G. Asphyxie périnatale et infirmité motrice d'origine cérébrale (I- Le diagnostic). *Gynécologie Obstétrique Fertil.* 2010. 1–38 p.
2. Everett TR, Peebles DM. Antenatal tests of fetal wellbeing. *Semin Fetal Neonatal Med.* juin 2015;20(3):138-43.
3. Carbonne B ME Pons K. Foetal scalp blood sampling during labour for pH and lactate measurements. *Best Pract Res Clin Obstet Gynaecol.* Vol. 30. 2016. 62–7 p.
4. Belfort MA S. GR. ST segment analysis as an adjunct to electronic fetal monitoring, Part I: background, physiology, and interpretation. *Clin Perinatol.* 1(143–157).
5. Ugwumadu A. Are we (mis)guided by current guidelines on intrapartum fetal heart rate monitoring? Case for a more physiological approach to interpretation. *BJOG Int J Obstet Gynaecol.* aout 2014;121(9):1063-70.
6. 15. Pinas A CE. Continuous cardiotocography during labour: Analysis, classification and management. *Best Pract Res Clin Obstet Gynaecol.* Vol. 30. 2016. 33–47 p.
7. Ayres-de-Campos D, Spong CY, Chandrharan E, FIGO Intrapartum Fetal Monitoring Expert Consensus Panel. FIGO consensus guidelines on intrapartum fetal monitoring: Cardiotocography. *Int J Gynaecol Obstet Off Organ Int Fed Gynaecol Obstet.* oct 2015;131(1):13-24.
8. Belfort MA, Saade GR, Thom E, Blackwell SC, Reddy UM, Thorp JM, et al. A Randomized Trial of Intrapartum Fetal ECG ST-Segment Analysis. *N Engl J Med.* 13 aout 2015;373(7):632-41.

9. Chandraharan E. Fetal scalp blood sampling should be abandoned: FOR: FBS does not fulfil the principle of first do no harm. *BJOG Int J Obstet Gynaecol.* 2016;123–11.
10. Classification CNGOF du rythme cardiaque fœtal : obstétriciens et sages-femmes au tableau ! *J Gynecologie Obstetrique Biol Reprod.* 2013;42–6.
11. Ayres-de-Campos D, Arteiro D, Costa-Santos C, Bernardes J. Knowledge of adverse neonatal outcome alters clinicians' interpretation of the intrapartum cardiotocograph. *BJOG Int J Obstet Gynaecol.* juill 2011;118(8):978-84.
12. Garabedian 3. C, De Jonckheere J, Butruille L, Deruelle P, Storme L, Houfflin-Debarge V. Understanding fetal physiology and second line monitoring during labor. *J Gynecol Obstet Hum Reprod.* 2017;46–2.
13. Chandraharan E. *Handbook of CTG Interpretation: From Patterns to Physiology.* 1^{re} ed. Cambridge, United Kingdom ; New York: Cambridge University Press; 2017. 256 p.
14. Ghesquiere L, De Jonckheere J, Drumez E, Sharma D, Aubry E, Deruelle P, et al. Parasympathetic nervous system response to acidosis: Evaluation in an experimental fetal sheep model. *Acta Obstet Gynecol Scand.* avr 2019;98(4):433-9.
15. Garabedian C, Champion C, Servan-Schreiber E, Butruille L, Aubry E, Sharma D, et al. A new analysis of heart rate variability in the assessment of fetal parasympathetic activity: An experimental study in a fetal sheep model. *PloS One.* 2017;12(7):e0180653.
16. Thacker SB, Stroup D, Chang M. Continuous electronic heart rate monitoring for fetal assessment during labor. *Cochrane Database Syst Rev.* 2001;(2):CD000063.
17. Alfirevic Z, Devane D, Gyte GM, Cuthbert A. Continuous cardiotocography (CTG) as a form of electronic fetal monitoring (EFM) for fetal assessment during labour. *Cochrane Database Syst Rev.* 03 2017;2:CD006066.
18. Jonsson M, Norden-Lindeberg S, Ostlund I, Hanson U. Metabolic acidosis at birth and suboptimal care—illustration of the gap between knowledge and clinical practice. *BJOG Int J Obstet Gynaecol.* oct 2009;116(11):1453-60.
19. Clinical negligence scheme for trusts. *Clinical risk management standards for maternity services.* London: NHS litigation Authority 2002.
20. Berglund S, Pettersson H, Cnattingius S, Grunewald C. How often is a low Apgar score the result of substandard care during labour? *BJOG Int J Obstet Gynaecol.* 1 juill 2010;117(8):968-78.
21. Jonsson M, Norden SL, Hanson U. Analysis of malpractice claims with a focus on oxytocin use in labour. *Acta Obstet Gynecol Scand.* 2007;86(3):315-9.
22. Nocon JJ, Coolman DA. Perinatal malpractice. Risks and prevention. *J Reprod Med.* fevr 1987;32(2):83-90.
23. Draycott T, Sibanda T, Owen L, Akande V, Winter C, Reading S, et al. Does training in obstetric emergencies improve neonatal outcome? *BJOG Int J Obstet Gynaecol.* fevr 2006;113(2):177-82.
24. Thellesen L, Bergholt T, Sorensen JL, Rosthoej S, Hvidman L, Eskenazi B, et al. The impact of a national cardiotocography education program on neonatal and maternal outcomes: A historical cohort study. *Acta Obstet Gynecol Scand.* 29 mai 2019;
25. Nielsen PV, Stigsby B, Nickelsen C, Nim J. Intra- and inter-observer variability in the assessment of intrapartum cardiotocograms. *Acta Obstet Gynecol Scand.* 1987;66(5):421-4.
26. Beaulieu MD, Fabia J, Leduc B, Brisson J, Bastide A, Blouin D, et al. The reproducibility of intrapartum cardiotocogram assessments. *Can Med Assoc J.* 1 aout 1982;127(3):214-6.

27. Palomaki O, Luukkaala T, Luoto R, Tuimala R. Intrapartum cardiotocography – the dilemma of interpretational variation. *J Perinat Med*. 2006;34(4):298-302.
28. Westerhuis MEMH, van Horen E, Kwee A, van der Tweel I, Visser GHA, Moons KGM. Inter- and intra-observer agreement of intrapartum ST analysis of the fetal electrocardiogram in women monitored by STAN. *BJOG Int J Obstet Gynaecol*. mars 2009;116(4):545-51.
29. Pehrson C, Sorensen JL, Amer-Wahlin I. Evaluation and impact of cardiotocography training programmes: a systematic review. *BJOG Int J Obstet Gynaecol*. juill 2011;118(8):926-35.
30. Morel O, Richard F, Thiebaugeorges O, Malartic C, Clement D, Akerman G, et al. [Fetal scalp pH: practical value during labour]. *Gynecol Obstet Fertil*. nov 2007;35(11):1148-54.
31. Stener Jorgensen. Fetal scalp blood sampling should be abandoned: AGAINST: Fetal scalp blood sampling in conjunction with electronic fetal monitoring reduces the risk of unnecessary operative delivery. *BJOG Int J Obstet Gynaecol*. 2016:123–11.
32. Reif P, Schott S, Boyon C, Richter J, Kavšek G, Timoh KN, et al. Does knowledge of fetal outcome influence the interpretation of intrapartum cardiotocography and subsequent clinical management? A multicentre European study. *BJOG Int J Obstet Gynaecol*. 2016;123(13):2208-17.

Hosted file

Figure 1.docx available at <https://authorea.com/users/299207/articles/428676-evaluation-and-impact-of-fetal-physiology-training-on-fetal-heart-rate-analysis>

Hosted file

Figure 2.docx available at <https://authorea.com/users/299207/articles/428676-evaluation-and-impact-of-fetal-physiology-training-on-fetal-heart-rate-analysis>