

Temperature-driven colour lightness and body size variation scale to local assemblages of European Odonata but are modified by propensity for dispersal

Daniel Acquah - Lamptey¹, Martin Braendle¹, Roland Brandl¹, and Stefan Pinkert¹

¹Philipps-Universität Marburg

April 28, 2020

Abstract

1. Previous macrophysiological studies suggested that temperature-driven colour lightness and body size variations strongly influence biogeographical patterns in ectotherms, but whether these trait-environment relationships scale to local assemblages and the extent to which they can be modified by dispersal remains largely unexplored. We test whether the predictions of the thermal melanism hypothesis and the Bergmann's rule hold for local assemblages. We also assess whether these trait-environment relationships are more important for species adapted to less stable (lentic) habitats, due to their greater dispersal propensity compared to those adapted to stable (lotic) habitats. 2. We quantified the colour lightness and body volume of 99 European dragon- and damselflies (Odonata) and combined these trait information with survey data for 518 local assemblages across Europe. Based on this continent-wide yet spatially explicit dataset, we tested for effects temperature and precipitation on the colour lightness and body volume of local assemblages and assessed differences in their relative importance and strength between lentic and lotic assemblages, while accounting for spatial and phylogenetic autocorrelation. 3. The colour lightness of assemblages of odonates increased and body size decreased with increasing temperature. Trait-environment relationships in the average and phylogenetic predicted component were equally important for assemblages of both habitat types but were stronger in lentic assemblages when accounting for phylogenetic autocorrelation. 4. Our results show that the mechanism underlying colour lightness and body size variations scale to local assemblages, indicating their general importance. These mechanisms were of equal evolutionary significance for lentic and lotic species, but higher dispersal ability seems to enable lentic species to cope better with historical climatic changes. The documented differences between lentic and lotic assemblages also highlight the importance of integrating interactions of thermal adaptations with proxies of the dispersal ability of species into trait-based models, for improving our understanding of climate-driven biological responses.

KEYWORDS

Bergmann's rule, dispersal, freshwater insects, habitat-stability-dispersal hypothesis, local assemblages, macrophysiology, Odonata, thermal adaptation, thermal melanism hypothesis

1 | INTRODUCTION

Understanding the processes that shape species' distributions and the composition of assemblages is within the centre of ecological research (Brown & Maurer, 1987; Cavender-Bares, Kozak, Fine, & Kembel, 2009; McGill, Enquist, Weiher, & Westoby, 2006; Ricklefs, 2004). A straightforward approach to a process-based understanding is to investigate functional traits that link the physiology of a species with the ambient environment in which the species occurs (Chown, Gaston, & Robinson, 2004; Violle et al., 2007). Ectothermic species must absorb thermal energy from their environment to be active and to maintain fundamental physiological processes, including growth and reproduction (Angilletta, 2009; Huey & Kingsolver, 1989). Therefore, ectotherms have evolved several behavioural (e.g. wing-whirring or basking; Corbet, 1980; May, 1979) and morphological adaptations to the climate in which they live (May, 1976; Angilletta, 2009).

Two of the most important morphological traits that influence the distribution of ectothermic organisms are probably their surface colour, particularly colour lightness (melanism), and body size. Fundamental physical principles link both traits to the heat gain and loss of an organism (Clusella-Trullas, van Wyk, & Spotila, 2007; Shelomi, 2012). On the one hand, melanisation of the cuticle determines the absorption of solar radiation and hence heat gain, a mechanism referred to as thermal melanism (Clusella-Trullas et al., 2007; Gates, 1980; Kalmus, 1941). On the other hand, since an increase in body size implies a reduction of the surface area to volume ratio, larger bodies are able to retain heat more efficiently than smaller bodies (Shelomi, 2012). Besides thermoregulation, greater melanisation increases resistance against pathogens, by enhancing the structural integrity of cells (Gloger's rule, Rapoport, 1969; Wilson, Cotter, Reeson, & Pell, 2001) and a larger body size is advantageous under dry conditions, as a lower surface area to volume ratio reduces water loss (Kühnel, Brückner, Schmelzle, Heethoff, & Blüthgen, 2017; R Emmert, 1981).

While the colour lightness and body size of a species should determine the climate in which it can live, the extent to which that species realises the potential environmental niche depends on its dispersal ability. The most important differences in species' abilities and propensities to disperse are related to the stability of their respective habitats. In general, species restricted to spatially and temporally stable habitats have lower dispersal abilities than species adapted to less stable habitats (Southwood, 1977). Freshwaters provide an ideal model system to test the predictions of this "habitat-stability-dispersal hypothesis" (Southwood, 1977). In the northern hemisphere, lentic water bodies (e.g., ditches and lakes) are ephemeral and date back to the Pleistocene, whereas the locations of rivers and streams (lotic waters) that carry water throughout the year have remained largely unaltered since the Mesozoic (Bohle, 1995 and sources therein). Species adapted to lentic waters have therefore evolved a suite of adaptations (i.e., trait syndromes) to cope with climatic changes, including morphological adaptations that facilitate mobility (Arribas et al., 2012; Hof, Brändle, & Brandl, 2006; Marten, Brändle, & Brandl, 2006; Pinkert et al., 2018) as well as behavioural adaptations (Corbet, 1980).

Recent studies have shown that the ecological differences between species adapted to lentic and lotic habitats carry a phylogenetic signal (Letsch, Gottsberger, & Ware, 2016). Moreover, these differences have led to contrasting biogeographical and diversification patterns between the two groups (Abellán, Millán, & Ribera, 2009; Hof, Brändle, & Brandl, 2008). For instance, Dehling et al. (2010) showed that the richness of lotic animals decreases from southern to northern Europe, whereas the richness of lentic animals is highest in central Europe. A broadly similar pattern has been reported for the richness of lentic and lotic Odonata (dragonflies and damselflies) on a global scale (Kalkman et al., 2008). Thus, in contrast to almost all other Odonata, the two youngest families (Coenagrionidae and Libellulidae; Rehn, 2003) that constitute the majority of lentic species globally (Kalkman et al., 2008) are disproportionately diverse in temperate climates. This suggests stronger trait-environment relationships in odonates of lentic than lotic habitats due to the greater ability of the former to cope with climatic changes in the past. However, despite strong theoretical reasons for an impact of species' dispersal ability on biogeographical patterns, to what extent dispersal can modify trait-environment relationships remains largely unexplored.

Analyses of the large-scale patterns of interspecific variation in physiological traits offer a powerful approach to elucidate the general processes that shape biodiversity patterns (Chown et al., 2004). These macrophysiological inferences are based on the assumption that the explanations for large-scale diversity patterns are found at lower levels of biological organisation, as functional traits influence the fundamental physiological rates of individuals and populations whereas the consequences thereof play an important role in determining a species' fundamental niche (Gaston & Blackburn, 2000). However, almost all of the studies conducted so far on the interspecific variation of colour lightness and body size in ectothermic species are based on expert range maps generated by interpolating species occurrence records across suitable habitats (e.g., Pinkert, Brandl, & Zeuss, 2017; Zeuss, Brandl, Brändle, Rahbek, & Brunzel, 2014; Zeuss, Brunzel, & Brandl, 2017). Hence, previous evidence of colour- and size-based thermoregulation has three important limitations. First, although at geographical scales expert range maps are generally considered to allow robust estimations of the full environmental range of species, the underlying distribution information tends to overestimate species' real distributional ranges (Hurlbert & Jetz, 2007; Merow, Wilson, & Jetz, 2017). Second, the inherent

spatial structure of expert range maps has been shown to inadvertently generate spurious spatial patterns for the richness and mean trait values of assemblages (Hawkins et al., 2017). Third, distribution data with a coarse resolution generate “synthetic” assemblages of species that do not necessarily form local assemblages. For instance, expert range maps typically also include records of populations that may no longer exist (or never existed) and pool species from different habitat types. Therefore, whether the previously documented relationships of colour lightness and body size with climate also scale to the local assemblage level remains largely unexplored.

In this study, we investigated trait-environment relationships using spatially explicit survey data for local assemblages of dragon- and damselflies (Odonata) across Europe. Specifically, according to the thermal melanism hypothesis and Bergmann’s rule *sensu lato*, we expected 1) an increase in the colour lightness of local assemblages of odonates with increasing temperature and 2) an increase in the body size of local assemblages of odonates with decreasing temperature. In addition, given that adaptations to spatially and temporally less stable habitats allow lentic species to better cope with climatic changes (habitat-stability-dispersal hypothesis), we predicted that the slopes of these relationships would be stronger for lentic than for lotic assemblages.

2 | MATERIAL AND METHODS

2.1 | Distribution data

Information on water body location and type (i.e. lentic or lotic) and the composition of local assemblages of odonates across Europe were compiled from data obtained in an extensive literature survey (Appendix 1). Only records of breeding species were included, to obtain species sets associated with the considered water bodies. Breeding records covered tandem pairs, ovipositing females, larvae, exuviae and recently emerged adults (Bried, Dillon, Hager, Patten, & Luttbeg, 2015), resulting in 5,703 records of 99 species of odonates and 524 local assemblages across 28 European countries. After assemblages with less than three species were excluded (to obtain reliable estimates of assemblage means), the final dataset comprised of 518 local assemblages of dragon- and damselflies (Fig. 1; 337 lentic, 181 lotic).

2.2 | Trait data

Following the most common approach used in the analysis of morphological traits based on digital images (Zeuss et al., 2014, 2017; Pinkert et al., 2017), we calculated the average colour lightness and body volume of species using drawings of European Odonata (Dijkstra & Lewington, 2006). To prepare images for the analysis, the body (head, abdomen, and thorax) in scanned drawings of species’ dorsal body surfaces (24-bits, sRGB, 1200 dpi resolution) was cropped out and saved to separate files using functions of Adobe Photoshop CS2. Based on these images the average colour of the pixels of an image across the red, green and blue channels was calculated as an estimate of the colour lightness of a species (8-bit grey values ranging from 0: absolute black to 255: pure white). In addition, these images were scaled with the magnification factor provided in Dijkstra and Lewington (2006) and used to calculate body volume in cm^3 ($\pi \times [\frac{1}{2} \text{ length of pixel row}]^2 \times \text{pixel edge length}$) as an estimate of the body size of a species based on the assumption that odonates generally have a cylindrical body form. The calculations were performed using functions of the R-package *png* (Urbanek, 2013). Body volume instead of linear size measures, such as wing length, head width and body length, was used because as a three-dimensional measure it allows for a more realistic estimate of the body mass of a species (Kühnel et al., 2017). Note that previous studies showed that the colour lightness and body volume estimates are correlated between drawings from different sources and between males and females (Pinkert et al., 2017, Zeuss et al., 2017). Subsequently, the average colour lightness and body volume were calculated across the species of each local assemblage. Previous studies found that the colour lightness and body size, as well as the habitat preference of European odonates, carries a phylogenetic signal (Zeuss et al. 2014; Letsch et al., 2016; Pinkert et al., 2017). Based on a recent phylogeny of the European odonates (Pinkert et al., 2018), we, therefore, partitioned the total variance of the colour lightness and body volume into a phylogenetic component and a species-specific component using Lynch’s comparative method (Lynch, 1991), implemented in the R-package *ape* (Paradis et al., 2004). The phylogenetic component

represents the variation in colour lightness and body volume predicted by the phylogenetic relationships of the species, whereas the species-specific component is the difference of the observed trait estimate from the phylogenetically predicted part. The advantage of this method is that it allows assessing the effect of phylogenetic signals in traits (i.e., P - component) that is often neglected as a source of bias, in addition to the model results that have been corrected for phylogenetic autocorrelation (i.e., S - component).

2.3 | Environmental data

Based on the predictions of the thermal melanism hypothesis and Bergmann's rule, we considered annual mean temperature as a predictor of geographical patterns in the colour lightness and body size of the assemblages of odonates. In addition, annual precipitation (AP) was included as a predictor because, in insects, lower surface-area to volume ratios of larger species are hypothesised to be advantageous under dry climate due to the associated reduction of water loss through the cuticle (Kühnel et al., 2017; Rimmert, 1981) and because of the protective function of melanin against pathogens under humid climates (e.g., Delhey, 2019; Rapoport, 1969; Stelbrink et al., 2019). Note that a principal component analysis based on the correlation of all 19 commonly used bioclimatic variables indicated that each of the two selected variables strongly contributed to the main annual trend in temperature and precipitation (Table S1). We considered only annual mean temperature and annual precipitation, rather than all 19 commonly used bioclimatic variables, to facilitate interpretations of their effects and comparability with other studies (e.g., Pinkert et al., 2017; Zeuss et al., 2017). Climate variables used in the analysis were extracted from climate data with a resolution of 2.5 arcminutes (retrieved from www.chelsea-climate.org; Karger et al., 2017, 2018), based on the geographical coordinates of the assemblages included in our analyses (with a buffer radius of approximately 1 km).

2.4 | Statistical analyses

All statistical analyses and calculations were conducted in R (version 3.5.1, R Core Team, 2016). Tests for trait-environment relationships were performed using single and multiple ordinary least-squares regression models, with the average colour lightness and body size of Odonata assemblages as dependent variables and climatic variables as independent variables. Differences in the slopes of the relationships of colour lightness and body size with climatic variables between lentic and lotic habitats were determined by fitting interaction terms between the independent variables and habitat type. In all models, independent variables were scaled and centred (z-standardised) to facilitate their comparison. To ensure very low multicollinearity among predictors, the variance-inflation factors (vif) for the predictor variables in regression models were checked using the *vif* function of the R-package *car* (Table S2; Fox et al., 2016).

Since spatial autocorrelation in the survey data could violate the assumptions of our statistical models, i.e., that all data points are independent of each other, spatial correlograms of the model residuals were calculated using functions of the R-package *ncf* (Bjørnstad, 2016). These correlograms indicated significant spatial autocorrelation in our data. Therefore, all analyses were repeated using spatial autoregressive error models (Dormann, 2007) that included a spatial distance weight according to the model-specific point of spatial independence (extracted from spatial correlograms shown in Figs S1-S2).

3 | RESULTS

3.1 | Trait-environment relationships

In all multiple regression models, the colour lightness of the assemblages of odonates increased with increasing annual mean temperature and body volume decreased with increasing annual mean temperature (Table 1). In all multiple regression models, the colour lightness of the assemblages of odonates was not affected by annual precipitation, but except for phylogenetically corrected models, body volume increased with increasing annual precipitation. These results were consistent with the results of single regression models (except that single regression of the average and phylogenetically predicted part of the variation in the colour lightness and annual mean temperature were not significant) and with the results of models that accounted for spatial autocorrelation (Table 1). The two climate predictors together explained up to between 20 % and 31 % of

the variation in colour lightness and between 2 % and 4 % of the variation in body volume (Table 1, see Table S3 for single regression models).

3.2 | Differences in trait-environment relationships between habitat types

The relative importance of annual mean temperature and annual precipitation, as well as the slopes of the considered trait-environment relationships, differed between lentic and lotic assemblages. Among the 24 considered relationships (models of the two traits, corrected for phylogenetic and spatial autocorrelation), 6 were significant for both lentic and lotic assemblages, 6 were only significant for lentic and 5 were only significant for lotic assemblages (Table 2). The two groups mostly responded similarly to annual mean temperature and all the relationships that were only significant in lotic assemblages were responses to annual precipitation (see Table S4 for single regression models).

Except for one response of body size, the slopes of all relationships of climate variables with the average as well as the phylogenetically predicted part of the variation in the traits were similar in their strength between the two groups. By contrast, responses of the species-specific part of the variation in colour lightness and body size to climate were mostly stronger in lentic compared to lotic assemblages (Table 2).

4 | DISCUSSION

Our results demonstrate that the spatial variation in colour lightness and body size (i.e., volume) of assemblages of odonates across Europe is mainly driven by temperature. In line with the predictions of the thermal melanism hypothesis and Bergman's rule *sensu lato*, our results showed that the analysed assemblages in warmer regions were consistently composed of, on average, lighter coloured and smaller species of dragon- and damselflies compared to assemblages in cooler regions. Our continent-wide yet spatially explicit assessment of these relationships reconciles previous macroecological (Pinkert et al., 2017; Zeuss et al., 2014, 2017) and experimental (e.g. May, 1991; Samejima & Tsubaki, 2010; reviewed in Clusella-Trullas et al., 2007) evidence indicating the general importance of mechanistic links of colour lightness and body size with the physiology and distribution of ectotherm species. In addition to the overall importance of colour- and size-based thermoregulation, our comparison of the trait-environment relationships of lentic and lotic assemblages of odonates revealed that the strength and relative importance of the climatic drivers of colour lightness and body size vary considerably between species with high and low dispersal/recolonisation ability.

Our study clearly showed that traits involved in thermoregulation influence the composition of dragon- and damselfly assemblages across Europe. According to the thermal melanism hypothesis, darker ectotherms are at an advantage in cool regions because of colour-based heat gain, and lighter ectotherms in warm regions because they reflect more solar radiation. In support of this hypothesis, we found that the colour lightness of Odonata assemblages increased with increasing temperature. The results of our analyses based on survey data together with the similar geographical patterns in colour lightness reported for assemblages of other ectothermic organisms at large geographical scales (Clusella-Trullas, Terblanche, Blackburn, & Chown, 2008; Heidrich et al., 2018; Schweiger & Beierkuhnlein, 2015; Stelbrink et al., 2019; Xing et al., 2018; Zeuss et al., 2014), confirm that thermal melanism is a mechanism of fundamental importance in ectothermic organisms across regions and scales. Furthermore, consistent with the predictions of Bergmann's rule *sensu lato*, we found that the average body size of assemblages of odonates decreased with increasing temperature. Even though a recent macroecological study by Zeuss et al. (2017) found support for Bergmann's rule in European odonates, its support in insects is generally equivocal (Shelomi, 2012), especially in studies conducted at small spatial and taxonomic scales. These contradictions in the results obtained at different scales have recently motivated debate about the reliability of large-scale assemblage-level studies, as it has been demonstrated that the type of distribution information on which most macroecological studies are based can purely by chance result in geographical patterns of species' traits (Hawkins et al., 2017). Despite temperature explained a comparatively low variance in body size (c.f. Zeuss et al., 2017), our findings support Bergmann's rule *sensu lato* in European odonates. Our support for both the thermal melanism hypothesis and Bergmann's rule using spatially explicit survey data for European odonates show that the findings of studies based on expert range maps are robust to pseudo-replications of co-occurrences and the inherent geographical structures of species

distributions (Hawkins et al., 2017).

Moreover, we also documented clear differences between species adapted to lentic and lotic habitats regarding the strength of the slopes of the considered trait-environment relationships and the relative importance of climatic drivers. Contrary to our third prediction, most of the relationships of average colour lightness and body size with temperature were equally strong between lentic and lotic assemblages. However, decomposing variations in colour lightness and body size showed that this is the result of similar responses of the phylogenetically predicted part of the traits of lentic and lotic species to climate, whereas relationships of the species-specific part of the traits were mostly stronger in lentic assemblages. Several studies have suggested that lentic species are stronger dispersers (e.g. Grewe, Hof, Dehling, Brandl, & Brändle, 2013; Hof et al., 2006; Marten et al., 2006) due to the negative relationship between habitat persistence and dispersal propensity (Southwood, 1962). Species adapted to lentic habitats are assumed to be closer to an equilibrium with ambient temperature (Dehling et al., 2010; Pinkert et al., 2018) and hence should dominate in recently recolonised regions (e.g., formerly glaciated northern parts of Europe; Pinkert et al., 2018). Accordingly, colour- and size-based thermoregulation together with high dispersal ability may have been hypothesised to cause contrasting biogeographical patterns between species adapted to lentic and lotic habitats over historical and evolutionary time scales (Hof et al., 2008; Pinkert et al., 2018). In fact, the distributional success and high diversity of lentic species in temperate regions seem to result not only from higher dispersal/recolonisation ability but also from an adaptive colour and body size evolution by lentic lineages. Our results suggest that adaptive colour and body-size are of similar importance for lentic and lotic species over evolutionary time scales, but that historical responses modified trait-environment relationships, with lentic species responding stronger to recent climatic changes than lotic species.

In light of previous zoogeographical and phylogeographical studies on dragon- and damselflies (Abellán et al., 2009; Kalkman et al., 2008; Pinkert et al., 2018; Sternberg, 1998), the documented differences in the trait-environment relationships of lentic and lotic species suggest that thermal melanism favours the colonisation of lineages of odonates in temperate climates. It has long been hypothesised that odonates are of tropical evolutionary origin and that only a few lineages acquired the ability to colonise and persist in temperate regions (e.g., Tillyard, 1916 p. 295). In a recent study, we found that the phylogenetic diversity of European Odonata assemblages decreased from the south-west to the north-east of the continent and that this pattern was mainly driven by the contemporary temperature (Pinkert et al., 2018). Latitudinal gradients of decreasing family or genus richness have been shown for odonates at the global scale (a simple proxy for the diversity of lineages; Kalkman et al., 2008). Furthermore, recent studies have documented a strong phylogenetic signal in the colour lightness of odonates and butterfly assemblages as well as differences in the importance of thermal melanism between butterfly families and associated these differences with a lower importance colour-based thermoregulation in tropical lineages (Zeuss et al., 2014, Pinkert et al., 2017, Stelbrink et al., 2019). Therefore, our finding that phylogenetically predicted part of the variation in colour lightness and body size is strongly driven by temperature suggested that colour- and size based thermoregulation might have played a central role in the adaptation to colder climates, whereas most Odonata lineages retained their initial tropical niche (see also Pinkert & Zeuss, 2018). Besides the differences in the strengths of the relationships of colour lightness and body size with temperature, our results show that the relative importance of temperature versus precipitation in shaping the geographical patterns of these traits differs between lentic and lotic assemblages. Although both annual mean temperature and annual precipitation consistently drove overall geographical patterns in the colour lightness and body volume of assemblages of odonates, lotic, but not lentic species seem to have an additional advantage of a higher size-based desiccation tolerance (Entling, Schmidt-Entling, Bacher, Brandl, & Nentwig, 2010), that also constrain their ability to thermoregulate via this trait. Specifically, we found that lotic assemblages in regions of lower precipitation were on average smaller than those in humid regions, which points to body size as an adaptation to water loss through the body surface (Kühnel et al., 2017). Furthermore, we showed that species adapted to lotic habitats were significantly larger in regions that are both warm and wet. This finding supports the predictions of Gloger’s rule (Wilson et al., 2001), which have been generally strongly supported by several large-scale studies (Pinkert et al., 2017; Stelbrink et al., 2019; Zeuss et al., 2014). Although studies have

shown that melanisation impacts desiccation resistance (Parkash, Rajpurohit & Ramniwas, 2008; Parkash, Sharma & Kalra, 2009), we are cautious about interpreting a potential colour-based protection against water loss for two reasons; firstly, the environmental gradient of the study sites did not include extreme humid or dry regions and secondly, in our study annual precipitation was not an important driver of the variations in colour lightness European Odonata assemblages.

5 | CONCLUSION

Our study highlights the importance of the mechanistic links of colour lightness and body size with the temperature regime which shapes the biogeographical patterns of dragon- and damselflies (Odonata). Colour- and size-based thermoregulation were by far the dominant mechanisms shaping the composition of assemblages of odonates, although other functions of body size and colour lightness seemed to influence the geographical patterns of both traits to some extent. The consistency of our findings together with the results of a number of macroecological analyses underlines the general importance of thermal melanism and Bergmann's rule for ectothermic organisms. However, besides highlighting the crucial role of traits involved in thermoregulation in shaping the distribution of odonate species, our results indicate that difference in species' dispersal propensities embedded in the spatio-temporal stability of their habitats contributes to explaining the scatter around the considered trait-environment relationships as well as to differences in the relative contributions of climatic predictors. Thus, thermal adaptations seem to be of similar evolutionary importance for lentic and lotic species but a greater dispersal ability of the former in combination with the climatic history of Europe seem to have allowed them better cope with historical climatic changes.

DATA AVAILABILITY STATEMENT

We intend to archive data for this study with Dryad Digital Repository.

ACKNOWLEDGEMENTS

Daniel Acquah - Lamptey acknowledges PhD funding by the Government of Ghana - Ministry of Education and the German Academic Exchange Service.

REFERENCES

- Abellán, P., Millán, A., & Ribera, I. (2009). Parallel habitat-driven differences in the phylogeographical structure of two independent lineages of Mediterranean saline water beetles. *Molecular Ecology* , 18(18), 3885-3902. <https://doi.org/10.1111/j.1365-294X.2009.04319.x>
- Angilletta, M. J. (2009). *Thermal Adaptation: A Theoretical and Empirical Synthesis* . Oxford, New York: Oxford University Press.
- Arribas, P., Velasco, J., Abellán, P., Sánchez-Fernández, D., Andújar, C., Calosi, P., ... Bilton, D. T. (2012). Dispersal ability rather than ecological tolerance drives differences in range size between lentic and lotic water beetles (Coleoptera: Hydrophilidae). *Journal of Biogeography* , 39(5), 984-994. <https://doi.org/10.1111/j.1365-2699.2011.02641.x>
- Bjornstad, O. N. (2016). ncf: Spatial Nonparametric Covariance Functions (Version R package version 1.1-7). Retrieved from Available at: <https://cran.r-project.org/package=ncf>
- Bohle, H. W. (1995). *Spezielle Ökologie: Limnische Systeme* . Berlin u.a.: Springer.
- Bried, J. T., Dillon, A. M., Hager, B. J., Patten, M. A., & Luttbeg, B. (2015). Criteria to infer local species residency in standardized adult dragonfly surveys. *Freshwater Science* , 34(3), 1105-1113. <https://doi.org/10.1086/682668>
- Brown, J. H., & Maurer, B. A. (1987). Evolution of species assemblages: Effects of energetic constraints and species dynamics on the diversification of the North American avifauna. *The American Naturalist* , 130(1), 1-17. <https://doi.org/10.1086/284694>

- Cavender-Bares, J., Kozak, K. H., Fine, P. V. A., & Kembel, S. W. (2009). The merging of community ecology and phylogenetic biology. *Ecology Letters* , 12(7), 693–715. <https://doi.org/10.1111/j.1461-0248.2009.01314.x>
- Chown, S. L., Gaston, K. J., & Robinson, D. (2004). Macrophysiology: Large-scale patterns in physiological traits and their ecological implications. *Functional Ecology* , 18, 159-167. <https://doi.org/10.1111/j.0269-8463.2004.00825.x>
- Clusella-Trullas, S., Terblanche, J. S., Blackburn, T. M., & Chown, S. L. (2008). Testing the thermal melanism hypothesis: a macrophysiological approach. *Functional Ecology* , 22, 232-238. <https://doi.org/10.1111/j.1365-2435.2007.01377.x>
- Clusella-Trullas, S., van Wyk, J. H., & Spotila, J. R. (2007). Thermal melanism in ectotherms. *Journal of Thermal Biology* , 32, 235-245. <https://doi.org/10.1016/j.jtherbio.2007.01.003>
- Corbet, P. S. (1980). Biology of Odonata. *Annual Review of Entomology* , 25 , 189-217.
- Dehling, D. M., Hof, C., Brändle, M., & Brandl, R. (2010). Habitat availability does not explain the species richness patterns of European lentic and lotic freshwater animals. *Journal of Biogeography* , 37(10), 1919-1926. <https://doi.org/10.1111/j.1365-2699.2010.02347.x>
- Delhey, K. (2019). A review of Gloger’s rule, an ecogeographical rule of color: Definitions, interpretations and evidence. *Biological Reviews* , 94, 1294-1316. <https://doi.org/10.1111/brv.12503>
- Dijkstra, K.-D. B., & Lewington, R. (2006). *Field Guide to the Dragonflies of Britain and Europe* . Dorset, UK: British Wildlife Publishing.
- Dormann, C. F. (2007). Effects of incorporating spatial autocorrelation into the analysis of species distribution data. *Global Ecology and Biogeography* , 16(2), 129-138. <https://doi.org/10.1111/j.1466-8238.2006.00279.x>
- Entling, W., Schmidt-Entling, M. H., Bacher, S., Brandl, R., & Nentwig, W. (2010). Body size–climate relationships of European spiders. *Journal of Biogeography* , 37(3), 477-485. <https://doi.org/10.1111/j.1365-2699.2009.02216.x>
- Fox, J., Weisberg, S., Adler, D., Bates, D., Baud-Bovy, G., Ellison, S., . . . Friendly, M. (2016). car: Companion to Applied Regression (Version R package version 2.1-4). Retrieved from Available at: <https://cran.r-project.org/package=car>
- Gaston, K. J., & Blackburn, T. M. (2000). *Pattern and process in macroecology* .
- Gates, D. M. (1980). *Biophysical Ecology* . New York: Springer.
- Grewe, Y., Hof, C., Dehling, D. M., Brandl, R., & Brändle, M. (2013). Recent range shifts of European dragonflies provide support for an inverse relationship between habitat predictability and dispersal. *Global Ecology and Biogeography* , 22(4), 403-409. <https://doi.org/10.1111/geb.12004>
- Hawkins, B. A., Leroy, B., Rodríguez, M. Á., Singer, A., Vilela, B., Villalobos, F., . . . Zelený, D. (2017). Structural bias in aggregated species-level variables driven by repeated species co-occurrences: a pervasive problem in community and assemblage data. *Journal of Biogeography* , 44(6), 1199-1211. <https://doi.org/10.1111/jbi.12953>
- Heidrich, L., Friess, N., Fiedler, K., Brändle, M., Hausmann, A., Brandl, R., & Zeuss, D. (2018). The dark side of Lepidoptera: Colour lightness of geometrid moths decreases with increasing latitude. *Global Ecology and Biogeography* , 27, 407-416. <https://doi.org/10.1111/geb.12703>
- Heinrich, B. (2013). *The Hot-Blooded Insects: Strategies and Mechanisms of Thermoregulation* . Springer Science & Business Media.
- Hof, C., Brändle, M., & Brandl, R. (2006). Lentic odonates have larger and more northern ranges than lotic species. *Journal of Biogeography* , 33(1), 63-70. <https://doi.org/10.1111/j.1365-2699.2005.01358.x>

- Hof, C., Brändle, M., & Brandl, R. (2008). Latitudinal variation of diversity in European freshwater animals is not concordant across habitat types. *Global Ecology and Biogeography* , 17(4), 539-546. <https://doi.org/10.1111/j.1466-8238.2008.00394.x>
- Huey, R. B., & Kingsolver, J. G. (1989). Evolution of thermal sensitivity of ectotherm performance. *Trends in Ecology & Evolution* , 4(5), 131-135. [https://doi.org/10.1016/0169-5347\(89\)90211-5](https://doi.org/10.1016/0169-5347(89)90211-5)
- Hurlbert, A. H., & Jetz, W. (2007). Species richness, hotspots, and the scale dependence of range maps in ecology and conservation. *Proceedings of the National Academy of Sciences* , 104(33), 13384-13389. <https://doi.org/10.1073/pnas.0704469104>
- Kalkman, V. J., Clausnitzer, V., Dijkstra, K. D. B., Orr, A. G., Paulson, D. R., & van Tol, J. (2008). Global diversity of dragonflies (Odonata) in freshwater. *Hydrobiologia* , 595, 351-363. <https://doi.org/10.1007/s10750-007-9029-x>
- Kalmus, H. (1941). Physiology and ecology of cuticle colour in insects. *Nature* , 148 , 428-431. <https://doi.org/10.1038/148428a>
- Karger, D. N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., ... Kessler, M. (2017). Climatologies at high resolution for the earth's land surface areas. *Scientific Data* , 4, 170122. <https://doi.org/10.1038/sdata.2017.>
- Karger, D. N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., ... Kessler, M. (2018). Data from: Climatologies at high resolution for the earth's land surface areas - Dryad Digital Repository. Retrieved July 9, 2018, <https://doi:10.5061/dryad.kd1d4>
- Kühnel, S., Brückner, A., Schmelzle, S., Heethoff, M., & Blüthgen, N. (2017). Surface area-volume ratios in insects. *Insect Science* , 24(5), 829-841. <https://doi.org/10.1111/1744-7917.12362>
- Letsch, H., Gottsberger, B., & Ware, J. L. (2016). Not going with the flow: a comprehensive time-calibrated phylogeny of dragonflies (Anisoptera: Odonata: Insecta) provides evidence for the role of lentic habitats on diversification. *Molecular Ecology* , 25(6), 1340-1353. <https://doi.org/10.1111/mec.13562>
- Lynch, M. (1991). Methods for the analysis of comparative data in evolutionary biology. *Evolution* 45, 1065-1080. <https://doi.org/10.1111/j.1558-5646.1991.tb04375.x>
- Marten, A., Brändle, M., & Brandl, R. (2006). Habitat type predicts genetic population differentiation in freshwater invertebrates. *Molecular Ecology* , 15(9), 2643-2651. <https://doi.org/10.1111/j.1365-294X.2006.02940.x>
- May, M. L. (1979). Insect Thermoregulation. *Annual Review of Entomology* , 24, 313-349. <https://doi.org/10.1146/annurev.en.>
- May, M. L. (1991). Thermal adaptations of dragonflies, revisited. *Advances in Odonatology* , 5 , 71-88.
- McGill, B. J., Enquist, B. J., Weiher, E., & Westoby, M. (2006). Rebuilding community ecology from functional traits. *Trends in Ecology & Evolution* , 21(4), 178-185. <https://doi.org/10.1016/j.tree.2006.02.002>
- Merow, C., Wilson, A. M., & Jetz, W. (2017). Integrating occurrence data and expert maps for improved species range predictions: Expert maps & point process models. *Global Ecology and Biogeography* , 26(2), 243-258. <https://doi.org/10.1111/geb.12539>
- Paradis, E., Claude, J. & Strimmer, K. (2004). APE: analyses of phylogenetics and evolution in R language. *Bioinformatics* , 20, 289-290. <https://doi.org/10.1093/bioinformatics/btg412>
- Parkash, R., Rajpurohit, S., & Ramniwas, S. (2008). Changes in body melanisation and desiccation resistance in highland vs. lowland populations of *D. melanogaster* . *Journal of Insect Physiology* , 54(6), 1050-1056. <https://doi.org/10.1016/j.jinsphys.2008.04.008>
- Parkash, R., Sharma, V., & Kalra, B. (2009). Impact of body melanisation on desiccation resistance in montane populations of *D. melanogaster* : Analysis of seasonal variation. *Journal of Insect Physiology* , 55(4), 898-908. <https://doi.org/10.1016/j.jinsphys.2009.06.004>

- Pinkert, S., Brandl, R., & Zeuss, D. (2017). Colour lightness of dragonfly assemblages across North America and Europe. *Ecography* , 40(9), 1110-1117. <https://doi.org/10.1111/ecog.02578>
- Pinkert, S., Dijkstra, K.-D. B., Zeuss, D., Reudenbach, C., Brandl, R., & Hof, C. (2018). Evolutionary processes, dispersal limitation and climatic history shape current diversity patterns of European dragonflies. *Ecography* , 41(5), 795-804. <https://doi.org/10.1111/ecog.03137>
- Pinkert, S., & Zeuss, D. (2018). Thermal biology: Melanin-based energy harvesting across the tree of life. *Current Biology* , 28(16), 887-889. <https://doi.org/10.1016/j.cub.2018.07.026>.
- Rapoport, E. H. (1969). Gloger's rule and pigmentation of Collembola. *Evolution* , 23(4), 622-626. <https://doi.org/10.1111/j.1558-5646.1969.tb03545.x>
- Rehn, A. C. (2003). Phylogenetic analysis of higher-level relationships of Odonata. *Systematic Entomology* , 28(2), 181-240. <https://doi.org/10.1046/j.1365-3113.2003.00210.x>
- Reilly, J. R., Hajek, A. E., Liebhold, A. M., & Plymale, R. (2014). Impact of *Entomophaga maimaiga* (Entomophthorales: Entomophthoraceae) on outbreak gypsy moth populations (Lepidoptera: Erebidae): The role of weather. *Environmental Entomology*, 43, 632-641. <https://doi.org/10.1603/EN13194>
- Remmert, H. (1981). Body size of terrestrial arthropods and biomass of their populations in relation to the abiotic parameters of their milieu. *Oecologia* 50, 12-13 <https://doi.org/10.1007/BF00378789>
- Ribera, I., Foster, G. N., & Vogler, A. P. (2003). Does habitat use explain large scale species richness patterns of aquatic beetles in Europe? *Ecography* , 26(2), 145-152. <https://doi.org/10.1034/j.1600-0587.2003.03271.x>
- Ricklefs, R. E. (2004). A comprehensive framework for global patterns in biodiversity. *Ecology Letters* , 7(1), 1-15. <https://doi.org/10.1046/j.1461-0248.2003.00554.x>
- Samejima, Y., & Tsubaki, Y. (2010). Body temperature and body size affect flight performance in a damselfly. *Behavioral Ecology and Sociobiology* , 64, 685-692. <https://doi.org/10.1007/s00265-009-0886-3>
- Schweiger, A. H., & Beierkuhnlein, C. (2015). Size dependency in colour patterns of Western Palearctic carabids. *Ecography* , 39(9), 846-857. <https://doi.org/10.1111/ecog.01570>
- Shelomi, M. (2012). Where are we now? Bergmann's rule sensu lato in insects. *The American Naturalist* , 180(4), 511-519. <https://doi.org/10.1086/667595>
- Southwood, T. R. E. (1962). Migration of terrestrial arthropods in relation to habitat. *Biological Reviews* , 37(2), 171-211. <https://doi.org/10.1111/j.1469-185X.1962.tb01609.x>
- Southwood, T. R. E. (1977). Habitat, the templet for ecological strategies? *Journal of Animal Ecology* , 46, 337-365.
- Stelbrink, P., Pinkert, S., Brunzel, S., Kerr, J., Wheat, C. W., Brandl, R., & Zeuss, D. (2019). Colour lightness of butterfly assemblages across North America and Europe. *Scientific Reports* , 9(1), 1760. <https://doi.org/10.1038/s41598-018-36761-x>
- Sternberg, K. (1998). The postglacial colonization of Central Europe by dragonflies, with special reference to southwestern Germany (Insecta, Odonata). *Journal of Biogeography* , 25, 319-337.
- Urbanek, S. (2013). png: Read and write PNG images (Version R package version 0.1-7). Retrieved from Available at: <https://cran.r-project.org/package=png>
- Violle, C., Navas, M.-L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I., & Garnier, E. (2007). Let the concept of trait be functional! *Oikos* , 116(5), 882-892. <https://doi.org/10.1111/j.0030-1299.2007.15559.x>
- Wilson, K., Cotter, S. C., Reeson, A. F., & Pell, J. K. (2001). Melanism and disease resistance in insects. *Ecology Letters* , 4(6), 637-649. <https://doi.org/10.1046/j.1461-0248.2001.00279.x>

Xing, S., Bonebrake, T. C., Ashton, L. A., Kitching, R. L., Cao, M., Sun, Z., . . . Nakamura, A. (2018). Colors of night: climate–morphology relationships of geometrid moths along spatial gradients in southwestern China. *Oecologia* , 188(2), 537-546. <https://doi.org/10.1007/s00442-018-4219-y>

Zeuss, D., Brandl, R., Brändle, M., Rahbek, C., & Brunzel, S. (2014). Global warming favours light-coloured insects in Europe. *Nature Communications* , 5, 4847. <https://doi.org/10.1038/ncomms4874>

Zeuss, D., Brunzel, S., & Brandl, R. (2017). Environmental drivers of voltinism and body size in insect assemblages across Europe. *Global Ecology and Biogeography* , 26, 154-165. <https://doi.org/10.1111/geb.12525>

BIOSKETCH

Daniel Acquah - Lamptey has a broad research interest in the ecology of insects, including thermal ecology, dispersal, and pest control. His research focuses on local scale interactions of Odonata assemblages with their biotic and abiotic environment.

Author contributions: D.A.-L., S.P. and R.B. conceived the ideas and designed methodology. D.A.-L. and S.P. collected the data. D.A.-L. and S.P. analysed the data. D.A.-L. wrote the first draft. S.P., R.B., and M.B. provided guidance, all authors contributed to revisions.

APPENDIX 1 - DATA SOURCES

Bang, C. Jr. (2001). Constructed wetlands: high-quality habitats for Odonata in cultivated landscapes, *International Journal of Odonatology* , 4(1), 1-15. <https://doi.org/10.1080/13887890.2001.9748153>

Bedjanic, M., & Salamun, A. (2003). Large golden-ringed dragonfly *Cordulegaster heros* Theischinger 1979, new for the fauna of Italy (Odonata: Cordulegastridae). *Natura Sloveniae* , 5(2), 19-29

Bernard, R. (2003). *Aeshna crenata* Hag., a new species for the fauna of Latvia (Anisoptera: Aeshnidae) *Notulae odonatologicae* , 6(1), 1-12.

Bogdanovic, T., Merdic, E., & Mikuska, J. (2008). Data to the dragonfly fauna of Lower Neretva River. *Entomologia Croatica* , 12(2), 51-65.

Bonifait, P. S., Defos Du Rau, P., & Soulet, D. (2008). Les Odonates de la Réserve Nationale de Chasse et de Faune Sauvage d'Orlu (département de l'Ariège, France). *Martinia* , 24(2), 35-44.

Bouwman, J., & Ketelaar, R. (2008). New records of *Coenagrion armatum* in Schleswig-Holstein (Odonata: Coenagrionidae). *Libellula* , 27(3/4), 185-190.

Braune, E., Richter, O., Söndgerath, D., & Suhling F. (2008). Voltinism flexibility of a riverine dragonfly along thermal gradients. *Global Change Biology* , 14, 470-482. <https://doi.org/10.1111/j.1365-2486.2007.01525.x>

Brochard, C., & van der Ploeg, E. (2013). Searching for exuviae of endemic Odonata species in Greece. *Brachytron* , 15(2), 83-101.

Buczyński P. (2006). General notes about the dragonfly (Odonata) fauna of the River Bug valley in the Lublin Region (SE Poland). In: R. Buchwald (Eds.), *Habitat selection, reproductive behaviour and conservation of central-European dragonflies (Odonata)*. *Proceedings of the 23rd Annual Meeting of the "Association of German-Speaking Odonatologists (GdÖ) at Oldenburg (Lower Saxony, Germany)* , (73-80). Aschenbeck & Isensee Universitätsverlag, Oldenburg.

Buczyński, P., & Tończyk, G. (2013). Dragonflies (Odonata) of Tuchola Forests (northern Poland). *Annales Universitatis Mariae Curie-Sklodowska , Section C Biologia* , 68 (1): 75-103. <https://doi.org/10.17951/c.2013.68.1.75>.

Buczyński, P. (1999). Dragonflies (Odonata) of sand pits in south-eastern Poland. *Acta hydrobiologia* , 41 (3/4), 219-230.

Buczyński, P., & Labeledzki, A. (2012). Landscape park of “Janowskie Forests” as a hotspot of dragonfly (Odonata) species diversity in Poland. In: D. H. Kazimierz (Eds.), *Natural human environment. Dangers,*

protection, education (pp. 151-174). University of Ecology and Management, Warsaw.

Buczyński, P., & Moroz, M. D. (2008). Notes on the occurrence of some Mediterranean dragonflies (Odonata) in Belarus. *Polish Journal of Entomology* , 77, 67-74.

Buczyński, P., Czachorowski, S., Moroz, M., & Stryjecki, R. (2003). Odonata, Trichoptera, Coleoptera, and Hydrachnidia of springs in Kazimierski Landscape Park (Eastern Poland) and factors affecting the characters of these ecosystems. *Supplementa ad Acta Hydrobiologica* , 5, 13-29.

Buczyński, P., Zawal, A., Stepień, E., Buczyńska, E., & Pešić, V. (2013). Contribution to the knowledge of dragonflies (Odonata) of Montenegro, with the first record of *Ophiogomphus cecilia*(Fourcroy, 1785). *Annales Universitatis Mariae Curie-Sklodowska, sectio C – Biologia*, 68(2), 57-71. <https://doi.org/10.17951/c.2013.68.2.57>.

Butler, S. G. (1993). Key to the larvae of European Orthetrum Newman (Anisoptera: Libellulidae). *Odonatologica*, 22(2), 191-196.

Carchini, G., Della Bella, V., Solimini, A. G., & Bazzanti, M. (2007). Relationships between the presence of odonate species and environmental characteristics in lowland ponds of central Italy. *Annales de Limnologie - International Journal of Limnology* , 43(2), 81-87. doi:10.1051/limn/2007020

Chavanec, A., & Raab, R. (2002). Die libellenfauna (Insecta: Odonata) des Tritonwassers auf der Donauinsel in Wien - ergebnisse einer langzeitstudie, aspekte der gewässerbewertung und der bioindikation. *Denisia* , 3, 63-79.

Chovanec, A., & Waringer, J. (2001). Ecological integrity of river–floodplain systems - assessment by dragonfly surveys (Insecta: Odonata). *Regulated Rivers: Research Management* , 17, 493-507. <https://doi.org/10.1002/rrr.664>

Clausnitzer, H-J. (1985). Die Arktische Smaragdlibelle (*Somatochlora arctica* ZETT) in der Südheide (Niedersachsen). *Libellula* , 4(1/2), 92-101.

De Knijf, G., & Demolder, H. (2010). Odonata records from Alentejo and Algarve, southern Portugal. *Libellula*, 29(1/2), 61-90.

Dolny, A., Drozd, P., Petříková, M., & Harabiš, F. (2010). Sex ratios at emergence in populations of some central European Gomphidae species (Anisoptera). *Odonatologica* , 39(3), 217-224.

Eggers, T. O., Grabow, K., Schutte C., & Suhling, F. (1996). Gomphid dragonflies (Odonata) in the southern tributaries of the river Aller, Niedersachsen, Germany. *Braunschweiger naturkundliche Schrriften*, 5(1), 21-34.

Ekestubbe, K. (1998). Artfördelning bland trollsländor (Odonata) i Södertälje kommun – analys av ett indikatorsystem för biologisk mångfald. *Södertälje kommun* , miljö- och.

Ellwanger, G., & Zirpel, S. (1995). Entwicklungsnachweis von *Anax imperator* Leach in einem Hochmoor in Estland (Anisoptera: Aeshnidae). *Libellula* , 14 (1/2), 41-48.

Farkas, A., Mérő, T. O., Móra, A., & Dévai, G. Y. (2014). Urban dragonflies: data on the odonata fauna of the Danube at Budapest. *Acta biologica Debrecina Supplementum oecologica Hungarica* , 32, 23-29.

Garcia, C. M. A. (1990). A key to the Iberian Orthetrum larvae (Anisoptera; Libellulidae). *Notulae odonatologicae*, 3(6), 81-96.

Golfieri, B., Hardersen, S., Maiolini, B., & Surian, N. (2016). Odonates as indicators of the ecological integrity of the river corridor: Development and application of the Odonate River Index (ORI) in northern Italy. *Ecological Indicators*, 61: 234-247. <https://doi.org/10.1016/j.ecolind.2015.09.022>.

Handke, K., Kalmund, P., & Didion, A. (1986). Die Libellen des Saarbrücker raumes. *Libellula* , 5(3/4), 95-112.

- Harabis, F., Tichanek, F., & Tropek, R. (2013). Dragonflies of freshwater pools in lignite spoil heaps: Restoration management, habitat structure and conservation value. *Ecological Engineering* , 55, 51-61. <https://doi.org/10.1016/j.ecoleng.2013.02.007>
- Hardersen, S. (2004). The dragonflies: species, phenology, larval habitats (Odonata). *Invertebrati Di Una Foresta Della Pianura Padana Bosco Della Fontana, Secondo Contributo – Conservazione Habitat Invertebrati* – 3, 29-50.
- Hardersen, S., & Dal Cortivo, M. (2008). Dragonflies (Odonata) of Vinchetto di Celarda Nature Reserve, with special reference to conservation actions. *Research on the Natural Heritage of Vinchetto Di Celarda and Val Tovanella (Belluno Province, Italy) – Quaderni Conservazione Habitat* – 5, 101-116.
- Hassal, C., Hollinshead, J., & Hull, A. (2012). Temporal dynamics of aquatic communities and implications for pond conservation. *Biodiversity and Conservation* , 21(3), 829-852. <https://doi.org/10.1007/s10531-011-0223-9>
- Heidemann, H., & Kull, R. (1986). Untersuchungen zur Libellenfauna und Gewässergüte an ausgewählten fließgewässern in Rheinland-Pfalz und Baden-Württemberg. *Libellula* , 5(1/2), 46-62.
- Holzinger, W. E. (1992). Die Libellenfauna der Mur-, Sulm- und Laßnitzauen des Leibnitzer Feldes (Steiermark, Österreich). *Libellula* , 11(3/4), 175-180.
- Horváth G. (2012). Assessment of riverine dragonflies (Odonata: Gomphidae) and the emergence behaviour of their larvae based on exuviae data on the reach of the river Tisza in Szeged. *Tiscia* 39, 9-15.
- Huber, A. (2000). On the odonate fauna of the Szamos (Someş) river and its surroundings in Romania. *Notulae odonatologicae* , 5(6), 69-84.
- Hübner, T. (1988). Zur Besiedlung neugeschaffener, kleiner Artenschutzgewässer durch Libellen. *Libellula* , 7(3/4), 129-145.
- Kalniņš, M. (2002). Banded Darter *Sympetrum pedemontanum*(Allioni, 1766) (Odonata: Libellulidae) – a new dragonfly species in the fauna of Latvia. *Latvijas Entomologs* , 39, 50-51.
- Kalniņš, M. (2006). The distribution and occurrence frequency of Gomphidae (Odonata: Gomphidae) in river Gauja. *Acta Universitatis Latviensis*, 710, 17-28.
- Keller, D., Seidl, I., Forrer, C., Home, R., & Holderegger, R. (2013). Schutz der Helm-Azurjungfer *Coenagrion mercuriale* (Odonata: Coenagrionidae) am Beispiel des Smaragd-Gebiets Oberaargau. *Entomo Helvetica* , 6, 87-99.
- Kitanova, D., Slavevska Stamenkovic, V., Kostov, V., & Marinov, M. (2008). Contribution to the knowledge of dragonfly fauna of the Bregalnitsa River, Macedonia (Insecta: Odonata). *Natura Montenegrina, Podgorica* , 7 (2), 169-180.
- Kolozsvári, I., Szabó, L. J., & Dévai, G. Y. (2015a). Occurrence pattern analysis of dragonflies (Odonata) on the river Tisza between Vilok and Huszt based on exuviae. *Applied Ecology and Environmental Research*, 13(4), 1183-1196. https://doi.org/10.15666/aeer/1304_11831196
- Kolozsvári, I., Szabó, L. J., & Dévai, G. Y. (2015b). Dragonfly assemblages in the upper parts of the river Tisza: a comparison of larval and exuvial data in three channel types. *Acta Zoologica Academiae Scientiarum Hungaricae*, 61(2), 189-204. <https://doi.org/10.17109/AZH.61.2.189.2015>.
- Koskinen, J. (2015). Dragonfly communities of north Karelian forest lakes and ponds. MSc. Thesis submitted to the Department of Biology of the University of Eastern Finland.
- Kulijer, D., Vinko, D., Billqvist, M., & Mekkes, J. J. (2012). Contribution to the knowledge of the Odonata fauna of Bosnia and Herzegovina – Results of the ECOO 2012. *Natura Sloveniae*, 14(2), 23-38.

- La Porta, G., Dell’Otto, A., Speziale, A., Goretti, E., Rebori, M., Piersanti, S., & Gaino, E., (2013). Odonata biodiversity in some protected areas of Umbria, central Italy. *Odonatologica* 42(2), 125-137.
- Macagno, A. L. M., Gobbi, M., & Lencioni, V. (2012). The occurrence of *Leucorrhinia pectoralis* (Charpentier, 1825) (Odonata, Libellulidae) in Trentino (Eastern Italian Alps). *Studi Trentini di Scienze Naturali*, 92, 33-36.
- Manenti, R., Siesa, M. E., & Ficetola, G. F. (2013). Odonata occurrence in caves: active or accidentals? A new case study. *Journal of Cave and Karst Studies* , 75(3), 205-209. <https://doi.org/10.4311/2012LSC0281>
- Marinov, M. (2007). Odonata of the Western Rhodopes, with special reference to the wetlands north of the town of Smolyan, South Bulgaria. *Notulae Odonatologicae* , 6(9), 97-108.
- Martens, A. (1991). Kolonisationserfolg von Libellen an einem neu angelegten Gewässer. *Libellula* , 10(1/2), 45-61.
- Mauersberger, R. (1990). Libellenbeobachtungen aus dem Bulgarischen Balkan-Gebirge (Stara Planina). *Libellula* , 9(1/2), 43-59.
- Moroz, M.D. (2013). Aquatic insects of cross-border water currents between Belarus and Ukraine. *Entomological Review* , 93(7), 874-886. <https://doi.org/10.1134/S0013873813070099>
- Moroz, M.D. Czachorowski, S. Lewandowski, K. & Buczyński, P. 2006. Aquatic insects (Insecta: Plecoptera, Ephemeroptera, Odonata, and Trichoptera) of the rivers in the Berezinskii Biosphere Reserve. *Entomological Review*, 86(9), 987-994. <https://doi.org/10.1134/S0013873806090028>
- Müller, Z., Jakab, T., Tóth, A., Dévai, G., Szállassy, N., Kiss, B., & Horváth, R. (2003). Effect of sports fisherman activities on dragonfly assemblages on a Hungarian river floodplain. *Biodiversity and Conservation*, 12, 167-179. <https://doi.org/10.1023/A:1021220220039>
- Muranyi, D., & Kovacs, T. (2013). Contribution to the Odonata fauna of Albania and Montenegro. *Folia Historico Naturalia Musei Matraensis* , 37, 29-41.
- Nelson, B., Ronayne, C., & Thompson, R. (2003). Colonization and changing status of four Odonata species, *Anax imperator*, *Anax parthenope*, *Aeshna mixta* and *Sympetrum fonscolombii* in Ireland 2000-2002. *The Irish Naturalists’ Journal* , 27(7), 266-272.
- Painter, D. 1998. Effects of ditch management patterns on odonata at Wicken fen, Cambridgeshire, UK. *Biological Conservation* , 84(2), 189-195. [https://doi.org/10.1016/S0006-3207\(97\)00095-5](https://doi.org/10.1016/S0006-3207(97)00095-5)
- Persson, S. (2011). Is the dragonfly composition changing in Central Sweden? BSc. thesis submitted to the Department of Nature conservation and species diversity of the Halmstad University, Finland.
- Piksa, K., Wachowicz, B., & Kwarciańska, M. (2006). Dragonflies (Odonata) of some small anthropogenic water bodies in Cracow City. *Fragmenta Faunistica*, 49(2), 81-89.
- Purse, B. V., & Thompson, D. J. (2003). Emergence of the damselflies, *Coenagrion mercuriale* and *Ceriagrion tenellum* (Odonata: Coenagrionidae), at their northern range margins, in Britain. *European Journal of Entomology* , 100, 93-99. <https://doi.org/10.14411/eje.2003.018>
- Raebel, E. M., Merckx, T., Riordan, P., Macdonald, D. W., Thompson, D. J. (2011). The dragonfly delusion: why it is essential to sample exuviae to avoid biased surveys. *Journal of Insect Conservation*, 14, 523-533. <https://doi.org/10.1007/s10841-010-9281-7>.
- Rajkov, S., Vinko, D., & Arandelovic, A. (2012). Faunistic results from the 2nd Balkan Odonatological Meeting – BOOM 2012, Serbia. *Natura Sloveniae* , 17 (2), 67-76.
- Rattu, A., Leo, P., Moratin, R., & Hardersen, S. (2014). *Diplacodes lefebvrei* in Sardinia, a new species for the Italian fauna (Odonata: Libellulidae). *Fragmenta entomologica* , 46(1-2), 121-124.

Reinhardt, K. (1997). Ein massenvorkommen mehrerer Libellenarten an einem Gewässer. *Libellula* , 16(3/4), 193-198.

Rozner, G., Lökkös A., & Ferincz, Á. (2010). Preliminary studies on the distribution of Large Golden Ringed Dragonfly (*Cordulegaster heros* Theischinger, 1979) and Golden Ringed Dragonfly (*Cordulegaster bidentata* Sélys, 1843) in the Kőszeg-mountains. *Folia Historico Naturalia Musei Matraensis* , 34, 37-40.

Rudolph, R. (1978). Notes on the dragonfly fauna of very small pools near Münster, Westfalia, German Federal Republic. *Notulae odonatologicae* , 1 (1), 1-16.

Sahlén, G. (1999). The impact of forestry on dragonfly diversity in central Sweden. *International Journal of Odonatology* , 2(2), 177-186. <https://doi.org/10.1080/13887890.1999.9748128>

Sahlén, G., & Ekestubbe, K. (2001). Identification of dragonflies (Odonata) as bioindicators of general species richness in boreal forest lakes. *Biodiversity and Conservation*, 10(5), 673-690. <https://doi.org/10.1023/A:1016681524097>

Schiel, F.-J. (1998). On the habitat requirements of *Erythromma najas* (HANSEMANN 1823) (Zygoptera: Coenagrionidae) in the Southern Upper Rhine valley. *Naturschutz südl. Oberrhein* , 2, 129-138.

Schindler, M., Fesl, C., & Chovanec, A. (2003). Dragonfly associations (Insecta: Odonata) in relation to habitat variables: a multivariate approach. *Hydrobiologia* , 497, 169-180. <https://doi.org/10.1023/A:1025476220081>

Schmidt, E. (1987). Zur Odonatenfauna des Geroldsees bei Garmlsch-Partenkirchen/Obb. (FRG) Ein Beitrag zur Analyse von Odonaten-Artenspektren bei kleiner Stichprobe. *Libellula* , 6(3/4), 121-134.

Schnapauff, I., Schridde, P., Suhling, F., & Ullmann, K. (1996). Libellenbeobachtungen in Nordost-Griechenland. *Libellula* , 15(3/4), 169-183.

Schridde, P. & Suhling, F. (1994). Larval dragonfly communities in different habitats of a Mediterranean running water system. *Advances in Odonatology* , 6, 89-100.

Schröter, A., Schneider, T., Schneider, E., Karjalainen, S., & Hämäläinen, M. (2012). Observations on adult *Somatochlora sahlbergi* – a species at risk due to regional climate change? (Odonata: Corduliidae). *Libellula* , 31(1/2), 41-60.

Schütte, C., & Suhling, F. (1997). Beobachtungen zum Fortpflanzungsverhalten von *Macromia splendens* (Pictet) (Anisoptera: Corduliidae). *Libellula* , 16(1/2), 81-84.

Thoma, M., & Althaus, S. (2015). Beobachtungen von Libellen (Odonata) auf dem Col de Bretolet (VS). *Entomo Helvetica* , 8, 97-109.

Weihrauch, F., & Weihrauch, S. (2006). Records of protected dragonflies from Rio Tera, Zamora province, Spain (Odonata). *Boletín Sociedad Entomológica Aragonesa* , 38, 337-338.

Wilderuth, H. (1992). Habitat selection and oviposition site recognition by the dragonfly *Aeshna juncea* (L.): an experimental approach in natural habitats (Anisoptera: Aeshnidae). *Odonatologica* , 22(1), 27-44.

Wittwer, T., Sahlén, G. & Suhling, F. (2010). Does one community shape the other? Dragonflies and fish in Swedish lakes. *Insect Conservation and Diversity* , 3, 124-133. <https://doi.org/10.1111/j.1752-4598.2010.00083.x>

TABLE 1 Effect sizes (z-scores) and the explained variance of predictor variables from single and multiple regressions (r^2/R^2) of the average, phylogenetic and species-specific components of the average colour lightness and body volume of 518 assemblages of European odonates with z-standardised temperature and precipitation variables. In addition, regression models (Nagelkerke pseudo- r^2/R^2) calculated with a spatial dependency weight are given. Significant relationships ($p < 0.05$) are shown in bold. The predictors are: annual mean temperature (AMT), and annual precipitation (AP). The P - component represents the phylogenetically predicted part of the respective trait and S - component represents the respective deviation of the average trait from the P - component.

Model	Model	Model	Trait	Predictor	Average	Average	S -	S -	P -
Ordinary	Ordinary	Single	Colour	AMT	Z-score	r^2/R^2	component	component	component
least-squares regression	least-squares regression		lightness		14.28	0.28	11.44	0.20	11.88
				AP	5.14	0.05	3.41	0.02	2.58
			Body volume	AMT	-1.59	0.01	-2.97	0.02	-1.33
		Multiple		AP	2.25	0.01	-0.25	0.00	2.55
			Colour lightness	AMT	13.12	0.29	10.80	0.20	11.55
				AP	1.68		0.33		-0.74
			Body volume	AMT	-2.40	0.02	-3.04	0.02	-2.21
				AP	2.89		0.68		3.11
Spatial autoregressive error models	Spatial autoregressive error models	Single	Colour lightness	AMT	11.62	0.30	9.87	0.21	10.73
				AP	2.69	0.14	1.47	0.09	1.17
			Body volume	AMT	-1.77	0.03	-2.50	0.02	-1.60
		Multiple		AP	2.24	0.03	0.77	0.01	2.45
			Colour lightness	AMT	11.32	0.31	9.57	0.21	10.64
				AP	1.71		0.50		-0.06
			Body volume	AMT	-2.15	0.04	-2.59	0.02	-2.00
				AP	2.56		1.04		2.74

TABLE 2 Individual slopes and standard error of the predictor variables from multiple regressions (R^2) of the average, phylogenetic and species-specific components of the average colour lightness and body volume of (lentic = 337, lotic = 181) assemblages of European odonates with z-standardised environmental variables. In addition, regression models (Nagelkerke pseudo- R^2) calculated with a spatial dependency weight are given. Shaded cells indicate significant differences in the slopes of these regressions between lotic and lentic assemblages. Slopes that are significant from zero ($p < 0.05$) are shown in bold. The predictors are: annual mean temperature (AMT), and annual precipitation (AP). The P - component represents the phylogenetically predicted part of the trait and S - component represents the respective deviation of the average trait from the P - component.

Model	Trait	Component	Predictor	Slope \pm SE for lentic	Slope \pm SE for lotic	R^2
Ordinary least-squares regression	Colour lightness	Average	AMT	$3.0 \times 10^0 \pm 2.8 \times 10^{-1}$	$2.7 \times 10^0 \pm 4.4 \times 10^{-1}$	0.29

Model	Trait	Component	Predictor	Slope \pm SE for lentic	Slope \pm SE for lotic	R^2
			AP	1.4×10^{-1} $\pm 2.9 \times 10^{-1}$	$8.6 \times 10^{-1} \pm 4.2 \times 10^{-1}$	
		S - component	AMT	1.2×10^0 $\pm 1.2 \times 10^{-1}$	6.6×10^{-1} $\pm 1.9 \times 10^{-1}$	0.21
			AP	$3.2 \times 10^{-2} \pm 1.3 \times 10^{-1}$	$2.6 \times 10^{-1} \pm 1.8 \times 10^{-1}$	
		P - component	AMT	1.6×10^0 $\pm 1.7 \times 10^{-1}$	1.7×10^0 $\pm 2.7 \times 10^{-1}$	0.22
			AP	$-9.7 \times 10^{-2} \pm 1.7 \times 10^{-1}$	$-1.5 \times 10^{-1} \pm 2.5 \times 10^{-1}$	
	Body volume	Average	AMT	$-2.5 \times 10^{-2} \pm 8.3 \times 10^{-3}$	$-1.2 \times 10^{-2} \pm 1.3 \times 10^{-2}$	0.03
			AP	$6.3 \times 10^{-3} \pm 8.5 \times 10^{-3}$	3.6×10^{-2} $\pm 1.2 \times 10^{-2}$	
		S - component	AMT	$-9.9 \times 10^{-4} \pm 6.4 \times 10^{-5}$	$-1.7 \times 10^{-4} \pm 1.0 \times 10^{-4}$	0.03
			AP	1.3×10^{-4} $\pm 6.5 \times 10^{-5}$	$-9.3 \times 10^{-5} \pm 9.5 \times 10^{-5}$	
		P - component	AMT	$-2.3 \times 10^{-2} \pm 8.3 \times 10^{-3}$	$-1.1 \times 10^{-2} \pm 1.3 \times 10^{-2}$	0.04
			AP	$6.7 \times 10^{-3} \pm 8.5 \times 10^{-3}$	3.9×10^{-2} $\pm 1.3 \times 10^{-2}$	
Spatial autoregressive error models	Colour lightness	Average	AMT	$2.9 \times 10^0 \pm 3.1 \times 10^{-1}$	$2.6 \times 10^0 \pm 4.6 \times 10^{-1}$	0.31
			AP	$3.0 \times 10^{-1} \pm 3.1 \times 10^{-1}$	$7.4 \times 10^{-1} \pm 4.3 \times 10^{-1}$	
		S - component	AMT	1.2×10^0 $\pm 1.3 \times 10^{-1}$	6.2×10^{-1} $\pm 2.0 \times 10^{-1}$	0.22
			AP	$7.8 \times 10^{-2} \pm 1.3 \times 10^{-1}$	$2.3 \times 10^{-1} \pm 1.9 \times 10^{-1}$	
		P - component	AMT	1.6×10^0 $\pm 1.8 \times 10^{-1}$	1.7×10^0 $\pm 2.7 \times 10^{-1}$	0.24
			AP	$4.1 \times 10^{-2} \pm 1.9 \times 10^{-1}$	$-1.5 \times 10^{-1} \pm 2.6 \times 10^{-1}$	

Model	Trait	Component	Predictor	Slope \pm SE for lentic	Slope \pm SE for lotic	R^2
	Body volume	Average	AMT	$-2.2 \times 10^{-2} \pm 8.8 \times 10^{-3}$	$-1.4 \times 10^{-2} \pm 1.3 \times 10^{-2}$	0.05
			AP	$8.1 \times 10^{-3} \pm 9.1 \times 10^{-3}$	$3.3 \times 10^{-2} \pm 1.3 \times 10^{-2}$	
		S - component	AMT	$-7.8 \times 10^{-5} \pm 6.8 \times 10^{-5}$	$-1.4 \times 10^{-4} \pm 1.0 \times 10^{-4}$	0.04
			AP	$1.8 \times 10^{-4} \pm 7.0 \times 10^{-5}$	$-9.1 \times 10^{-5} \pm 9.7 \times 10^{-5}$	
		P - component	AMT	$-2.2 \times 10^{-2} \pm 8.8 \times 10^{-3}$	$-1.4 \times 10^{-2} \pm 1.3 \times 10^{-2}$	0.05
			AP	$8.4 \times 10^{-3} \pm 9.2 \times 10^{-3}$	$3.6 \times 10^{-2} \pm 1.3 \times 10^{-2}$	




FIGURE 1 Distribution of (lentic = 337, lotic = 181) assemblages odonates across Europe. Lentic assemblages are indicated with red circles, lotic assemblages with blue circles and the main European rivers in grey. The black outlines are country boundaries in the study region. The proportion of lotic habitats is higher in southern and central Europe. The space inside each circle represents the area for which the climate data were aggregated (a radius of ~ 1 km around the location of each community).




FIGURE 2 Scatterplots of the average (a, b), species-specific (c, d) and phylogenetic (e, f) components of the average colour lightness and body volume of (all habitats = 518, lentic = 337, lotic = 181) assemblages of European odonates and z-standardised annual mean temperature, and annual precipitation. The colour of the dots indicates the habitat type of the assemblages. Lines indicate regression lines of ordinary least squares models. The colour lightness ranges from 0 (absolute black) to 255 (pure white). The P - component represents the phylogenetically predicted part of the trait and S - component represents the respective deviation of the average trait from the P - component.