

Collateral effects of COVID-19 pandemic in pediatric hemato-oncology: fatalities caused by diagnostic delay

Rosanna Parasole¹, Pio Stellato², Valentino Conter³, Antonia De Matteo², Luigia D'Amato², Antonella Colombini³, Carmine Pecoraro², Carmela Bencivenga², Marta Raimondo², Susanna Silvestri², Vincenzo Tipo², Luigi Annichiarico Petruzzelli², Giovanna Giagnuolo¹, Agostino Curatolo², Andrea Biondi³, and Giuseppe Menna⁴

¹Santobono-Pausilipon Hospital

²AORN Santobono Pausilipon

³Ospedale S. Gerardo, University of Milano-Bicocca, Fondazione MBBM, Monza, Italy

⁴Pausilipon Hospital

May 20, 2020

Letter to the Editor

Coronavirus disease COVID-19 has deeply modified national health services with a profound impact on hospital and in particular emergency and intensive care units (ICU) activities. As recently reported in Italy pediatric emergency accesses substantially decreased likely due to the instructions to prevent overcrowding in emergency rooms and spread of SARS-CoV-2 infection and to fear of the infection.¹ At the Santobono-Pausilipon Hospital (Naples), pediatric emergency accesses in March 2020 were only one fifth of those registered in 2019 in the same period. Likewise a marked reduction of consultations occurred also in family pediatricians clinics.²

We report here 3 children who arrived at hospital in life-threatening conditions at the onset of Acute Lymphoblastic Leukemia (ALL) between March 14 and April 10, 2020.

First case: a 2-year-old-child arrived at the emergency department with a 15 days history of fatigue, pallor and dyspnea, in a comatose state, with severe anemia, respiratory distress, hematemesis and metabolic acidosis. Chest X-ray showed interstitial pneumonia. Blood tests showed: hemoglobin 2.7 gr/dL, WBC count 185.000/ μ l, platelets (PTL) 10.000/ μ l, LDH 3609 U/L. Peripheral blood was diagnostic for CD10, CD19 and CD58 positive ALL (B-lineage ALL). The patient, admitted at the ICU, intubated, transfused with RBC, PTL and plasma, died 12 hours after arrival at the hospital due to progressive worsening of clinical conditions. The nasal swab was negative for SARS-CoV-2 and positive for adenovirus.

Second case: a 5-year-old-child arrived at the emergency department with a one month history of respiratory distress. Imaging showed a mediastinal mass compressing the brachiocephalic vein, the aorta, the pulmonary trunk and the left pulmonary artery, tracheal deviation, compression of the left main bronchus, left lung atelectasis and pleural effusion. Blood tests showed: hemoglobin 14.5 gr/dL, WBC count 37.000/ μ l, PTL 294.000/ μ l, LDH 6153 U/L, creatinine 1.9 mg/dl. Peripheral blood was diagnostic for CD5, CD7, CyCD3 and CD8 positive ALL (T-ALL). Steroid treatment was started. Clinical conditions deteriorated rapidly with cardiac and renal failure. The patient, admitted to ICU 2 hours after arrival at the hospital and intubated, died 24h later. The nasal swab was negative for SARS-CoV-2.

Third Case: a 4-year-old child arrived at the hospital with one month history of fever, cough and shortness of breath treated at home with antibiotics and steroids without improvement. Imaging showed a mediastinal

mass compressing the left brachiocephalic, azygos and superior cava veins, and right pulmonary artery and vein; mild tracheal deviation, compression of the left main bronchus; pericardial and pleural effusion; nephro-hepato-splenomegaly and ascites. Due to signs of cardiac tamponade, pericardiac and pleural drainage were placed and the patient was admitted at ICU and intubated. Blood tests showed: normal hemoglobin, WBC and PTL counts; LDH 2732 U/L, creatinine 2.98 mg/dl, K 8 mEq/L, Ca 5.4 mEq/L. Bone marrow was diagnostic for CD2, CD5, CD7, CD99 and CyCD3 positive ALL (T-ALL). Treatment with steroids was started. Due to progressive renal failure hemodialysis was performed for 9 days. Clinical conditions improved with rapid shrinking of mediastinal masses and resolution of pericardial and pleural effusion. The patient was thus extubated and treatment for ALL was instituted with good response to induction therapy. The nasal swab was negative for SARS-CoV-2.

The 3 cases of ALL here described, 2 of them fatal, arrived at the hospital in critical conditions, most likely as a consequence of fear of COVID-19. Delay in diagnosis of neoplastic disease is a well-known problem in low-middle income countries (LMIC), but is quite rare in high-income countries (HIC). Actually, this combination of events never occurred in the past at the Santobono-Pausilipon Hospital, where, at the time of writing, no SARS-CoV-2 positive cases have been identified among children treated for cancer.

Considering low prevalence of virus spreading in children and that SARS-CoV-2 positive children are generally asymptomatic or have a very mild course of the disease there is a substantial risk that collateral effects of COVID-19 pandemic, i.e. delays in diagnosis, chemotherapeutic treatments and treatment of chemotherapy complications, may be worse than those posed by the disease itself.^{3,4,7} Recently the major pediatric cancer scientific associations have expressed great concern on the risk that fear to access to medical care raised by Covid-19 may cause these delays not only in LMIC but also in HIC with dramatic consequences we are not used to face.⁵⁻⁶ Our experience confirms the occurrence of these collateral effects, indicating that there is a need of awareness of this risk and careful medical attention to assure timely diagnoses and adequate treatment adherence in childhood cancer.

Conflict of Interest Statement

No conflict of interest to disclose

References

1. ISS—COVID-19 Italian epidemiological report. <https://www.epicentro.iss.it/coronavirus/sars-cov-2-sorveglianza-dati>
2. Lazzerini M., et al. Delayed access or provision of care in Italy resulting from fear of covid 19. *Lancet Child Adolesc Health* 2020.[https://doi.org/10.1016/S2352-4642\(20\)30108-2](https://doi.org/10.1016/S2352-4642(20)30108-2)
3. Saini K.S., de las Heras B., de Castro J., et al. Effect of the COVID-19 pandemic on cancer treatment and research. *Lancet Haematol* 2020 [https://doi.org/10.1016/S2352-3026\(20\)30123-X](https://doi.org/10.1016/S2352-3026(20)30123-X)
4. Casanova M, Pagani Bagliacca E, Silva M, et al. How young patients with cancer perceive the COVID-19 (coronavirus) epidemic in Milan, Italy: Is there room for other fears? *Pediatr Blood Cancer* 2020.<https://doi.org/10.1002/pbc.28318>
5. Balduzzi A., Brivio E., Rovelli A., et al. Lessons after the early management of the COVID-19 outbreak in a pediatric transplant and hemato-oncology center embedded within a COVID-19 dedicated hospital in Lombardia, Italy. *Estote parati. Bone Marrow Transplantation*2020.<https://doi.org/10.1038/s41409-020-0895-4>
6. Sullivan M, Bouffet E, Rodriguez-Galindo C, et al. The COVID-19 PANDEMIC: A Rapid Global response for Children with Cancer from SIOP, COG, SIOP-E, SIOP-PODC, IPSO, PROS, CCI and St Jude Global. *Pandemic COVID-19 PBC Final* 27 April 2020
7. Hrusak O, Kalina T, Wolf J, Balduzzi A, et al. Flash Survey on Severe Acute Respiratory Syndrome coronavirus-2 Infections in Paediatric Patients on Anticancer Treatment. *Eur J Cancer.* 2020 Apr 7;132:11-16. doi: 10.1016/j.ejca.2020.03.021.