

Qualitative Olfactory Disorders: Patient experiences and self-management

Carl Philpott¹, Joanne Dixon², and Duncan Boak²

¹University of East Anglia

²Affiliation not available

June 22, 2020

Key points

Question

What are the specific experiences of olfactory disorder patients affected by the qualitative symptoms of parosmia and phantosmia including both triggers for symptoms and self-help measures they have tried?

Findings

Parosmia was reported as a daily symptom in 67% compared to 31% for phantosmia. Only 4% of respondents reported having received any successful treatment for their qualitative symptoms and 58% reported having received no treatment whatsoever. Stimulating the nose with other smells was the most common self-management method despite 61% stating external odours as triggers for their symptoms.

Meaning

The duration of qualitative olfactory symptoms, the relative lack of interest from medical professionals and the lack of therapeutic options for them mean there is a significant burden for these patients and further research for treatment options and coping strategies is needed.

Introduction

Background and rationale

Whilst we are now witnessing the rise of sudden onset anosmia as a marker of Covid-19 coronavirus infection¹⁻⁷, it may serve to remind us that at least an estimated 5% of the population⁸ are affected by olfactory dysfunction. These disorders can be both quantitative and qualitative in nature. Common causes of olfactory disorders include chronic rhinosinusitis, post-viral olfactory loss and post-traumatic olfactory loss⁹ with neurological diseases such as Parkinson's sometimes culpable, but with about 10% of all cases in the community appearing to be idiopathic in nature. Amongst those cases typically common in specialist clinics are post-viral and idiopathic cases¹⁰. The qualitative aspects of olfactory disorders in the form of parosmia and phantosmia are however less well publicised¹¹.

Fifth Sense, the UK charity for people affected by smell and taste disorders, was founded in 2012 and working together with patients our previous work has been able to demonstrate the very significant impact of olfactory

disorders on those affected^{12,13}. Whilst recent findings show that qualitative disorders may have a positive prognostic significance for the recovery of olfactory disorders¹⁴ they do however present a particular source of distress to those affected whilst present. Previous studies of patients presenting with olfactory dysfunction have shown that parosmia was present in 34% and phantosmia present in 12% of patients presenting to a clinical setting¹⁵. In some of our earlier work, 2 years after the inception of our charity, our membership reported parosmia in 19% and phantosmia in 24% of cases¹².

Even more so than quantitative olfactory loss, qualitative disturbances provide a source of frustration for both the patient and the clinician. Evidence for prescribed treatment for qualitative disorders was limited to 7 studies in a previous systematic review for phantosmia¹⁶ and evidence for self-help measures appears to be very limited¹⁷. Some patients are so severely affected that consideration has been given to surgical removal of the olfactory epithelium¹⁸.

Objectives

This study aimed to characterise the specific experiences of olfactory disorder patients affected by the qualitative symptoms of parosmia and phantosmia including both triggers for symptoms and self-help measures they have tried.

Materials and Methods

Study design

The study was designed as a cross-sectional survey of the personal experience of people affected by qualitative olfactory disorders in living with and managing these symptoms. A survey questionnaire was developed using a clinician-expert patient partnership. The survey ran online and was available for a period of 6 months. It was promoted via social media internationally using the Fifth Sense links. One of our charity's volunteers (JD) also provided a personal account of her experience of parosmia as well as inviting comments from her local support group in Newcastle.

Setting and Participants

The survey was open to anyone globally with access to the world wide web and declaring themselves as someone affected by the symptoms of parosmia and phantosmia, regardless of the aetiology. Survey participants were able to access the survey themselves free of charge via the web-based platform SurveyMonkey. Participants were self-selecting and could participate from any country internationally.

Data sources/management and variables

The survey asked for basic demographics including age and sex. Participants were asked to declare the underlying cause for their smell loss. Further questions explored the timing, intensity, frequency and impact of qualitative symptoms along with any triggers and self-management undertaken.

Bias

We aimed to reduce bias in responses by making the survey widely available and due to the membership base of charity, reflects both a UK population and an international membership.

Study size and Statistical Methods

As this was an exploratory study to capture patient experiences, no sample size was set. Due to the nature of the study, descriptive statistics only have been utilised in reporting the survey data.

Results

Participants

A total of 100 participants recorded information on the survey during the study period.

Descriptive data

Of the 100 participants, 61 were female and 39 were male. The age of participants ranged from 13 to 88, with a mean age of 51. The aetiology reported for participants included 33% reporting idiopathic olfactory loss (IOL), 17% chronic rhinosinusitis (CRS), 23% post-traumatic olfactory loss (PTOL) and 26% post-viral olfactory loss (PVOL) and 5% iatrogenic; 4 patients reported both PVOL and CRS. The range of duration of reported qualitative olfactory disorders was 1 month to 40 years with a mean of 7 years. Eighty percent of respondents reported experiencing with parosmia whilst 65% reported phantosmia.

Main results

Frequency and impact of symptoms

Symptoms of parosmia were reported as occurring more frequently (table 1), with it being reported as a daily symptom in 67% compared to 31% for phantosmia (table 2) with an associated higher quality of life impact. The onset of the symptoms was reported as being sudden in 65% and the duration of symptoms ranged from 1 month to over 40 years. Parosmia was more likely to be constant (64%) than phantosmia which was reported as fluctuating in 59%.

Self-management of symptoms

Only 4% of respondents reported having received any successful treatment for their qualitative symptoms and 58% reported having received no treatment whatsoever. Stimulating the nose with other smells was the most common self-management method (table 3), despite the fact that 61% described other smells as a trigger for their symptoms (table 4). Several reported the use of nasal douching, but other options left in the comments included blocking nasal passages with tissue or nasal plugs, using dried eucalyptus leaves in boiling water and inhaling the steam and drinking the tea inhaling the steam, thyme and lavender and castor oil drops in the nose. Comments on stimuli included the random or constant nature in some that was independent of any stimuli, with others reporting increased rhinitis symptoms, cooking smells are the worst, emotional or tense situations, humidity and temperature changes, exercise, memories and visual cues, eating and fatigue.

Personal account

Having suffered an extremely bad cold in 2015 (aged 44) I was unaware of the putrid bad smell and taste that was soon to follow and consume my daily life. Over the coming weeks this foul smell became so overpowering that sleep was my only escape. I started to notice certain smells i.e. coffee, petrol, smoke, cooked food, perfume, fabric softener intensified the smell of rotten flesh and sewage twentyfold. Washing clothes, using

soap, deodorant and shampoo were all revolting and using toothpaste would make me retch. It took over my life and affected my work and personal life in such a way that I was off work for 3 months and avoided being around my partner, going out or seeing friends and family. This bad smell was constant even when there were no other odours present. The only place I could stop the smell from intensifying was at home by avoiding certain triggers as mentioned above.

I saw my GP over several months who prescribed a variety of medication none of which worked. To my amazement one GP suggested I stand on my head to clear my sinuses. My nostrils became very dry and crusts developed alongside nose bleeds. Eventually I came across a local ENT consultant in the region who took an interest in these conditions and my GP made an appointment for me, but this would not be for another 6 months. Having explored so much medication without success I became so desperate that I arranged to see this consultant privately and was seen within days. Unfortunately, they confirmed that I was suffering from parosmia but could not tell me if I would regain my smell and taste and there was nothing more that could be done. At that point I became hysterical and totally broke down - he recommended I speak to my GP about antidepressants. I left his office distraught and feeling my life was over – I could not bear this constant repulsive smell and taste any longer. Thankfully a couple of days later my consultant called who'd spoke with a colleague and recommended I be prescribed with Theophylline. He explained there was no evidence to prove it would be successful as it was a trial drug which was used for asthma patients. After 1-2 weeks of taking this medication I found it had suppressed the bad smell to a degree where I could manage. I also noticed that mucus returned in my nose as things slowly started to improve but by this time I had a perforated septum by blowing my nose so hard and dislodging the crusting in order to help breathe. Having read about a case in the US regarding Gabapentin I decided to ask my GP if I could also be prescribed this medication and over time I felt that this helped improve my taste.

Parosmia is a dreadful condition which seriously impacted my mental health and quality of life. However, some people do recover – I have! It may have taken 5 years, but I can say I have 85% of my normal smell and taste back. Having spoken to other people with the same condition it is awful to hear that some GPs and Consultants are uninterested and clinical in telling patients "If your smell doesn't return after 6 months then it's not likely to." In my experience GPs/Consultants should be supportive and monitor and measure their patients' progress. They should be made aware of specific medication that can potentially aid recovery and save on cost spent on medication that will have no effect. There should also be more research done on trial drugs.

I fully understand that the medical profession is unable to prove that a patient can recover from parosmia but equally there aren't any statistics to say that a patient won't recover either. Therefore, taking an interest and having a positive approach will support patients on their potential journey to recovery or at least give them time to slowly adapt and find coping mechanisms.

Feedback from Fifth Sense Newcastle hub members

- Depression/anxiety when living alone, afraid to go out with friends for fear of eating something that suddenly causes a serious chemical taste/smell and I have to leave. The taste e.g. dressings with vinegar - lemon juice. Loss of joy of smelling cut grass, sea, flowers, personal contact. Life is so bland - feel alien.
- It is depressing that I might have to suffer with this for the rest of my life. My GP tells me that considering the health problems of most of his 75-year-old patients my problem is minor. I know it's not life threatening but it does seriously affect my quality of life.
- My consultant seemed spectacularly uninterested and spent just a few minutes with a cursory examination to then tell me that there was 'no hope' for me! Crushed, I all but gave up, and was then assaulted by a further issue where my no-taste was largely replaced with an unpleasant sensation/taste for most foods.
- I can't live with this condition - having this foul smell and taste every second of the day is just unbearable.

- *The condition manifests itself for me by fairly extreme and unpleasant smells/tastes - somewhat akin to the sharp, sour smell of vomit - for several types of foods (coffee, beef, eggs, broccoli etc) - but all those different foodstuffs have the SAME unpleasant odour!! Weird in the extreme! And...occasionally, things I could tolerate before, I suddenly become unable to? And, yes, like the leaflet indicates, I feel that I was "better off" when I couldn't smell or taste anything!*

Discussion

Key results

Qualitative olfactory disorders can have a significant impact on patients and a lasting effect given the mean duration of symptoms in this cohort was 7 years. A lack of treatment or effective treatment was evident in 96%. The combination of these two factors reveals a group of patients with a poor quality of life and the potential for adverse mental health. Many triggers exist but whilst external odours do stimulate symptoms, they are also a means for patients to try to moderate their symptoms.

Limitations

The survey will not have been seen by those who are not online or do not have access to the aforementioned social media. This is likely to have disproportionately affect the older generations. The charity membership and survey respondents will also tend to be more likely to be those who have more persistent symptoms and perhaps are more resistant cases. The diagnoses were self-reported and no physical examination or psychophysical olfactory testing was conducted due to the nature of the study setting. However, we believe the responses on the survey to genuinely reflect the experiences of our charity's members based on our previously published work ^{12,13} and the feedback during our members conferences and workshops ¹⁹.

Interpretation

The demographics and aetiology of study participants was in keeping with the typical female predominance seen in other studies but interestingly with idiopathic and CRS accounting for the aetiology as of many of the respondents as PVOL and PTOL. Previous studies have reported the prevalence of qualitative disorders as most frequent in patients with PVOL (56%), with 28% in CRS, 14% in PTOL and 10% in idiopathic cases ¹⁵. Recovery of qualitative olfactory dysfunction in the study by Reden et al was seen in 29% of those with parosmia after an average of 12 months and in 53% of those with phantosmia. A previous systematic review of the management of phantosmia specifically found a very small evidence base for prescribed treatments that included antipsychotics, antiepileptics, antimigrainous medications and topical cocaine ¹⁶. Clearly this group of patients have an unmet need in terms of effective treatments, given that 38% of our participants reported unsuccessful treatment and over half reported no treatment being given.

The mean duration of symptoms seen in our study was longer than otherwise reported ^{15,20} but may reflect the source of recruitment and therefore encapsulating patients whose symptoms persist beyond their immediate clinical contact. The main odorant triggers eliciting parosmia in a study by Pierre et al included petrol, tobacco, coffee, perfumes, fruits and chocolate ²⁰. Our volunteers encapsulate a range of experience and that the experience of parosmia is very much an individual one.

Generalisability

Qualitative disturbances remain problematic for those who experience them due to the duration of symptoms, the relative lack of interest from medical professionals and the lack of therapeutic options for them. This has a significant impact on their quality of life ²⁰. In future, consideration needs to be given to psychological and dietary measures to enable patients to adapt and recover from these disabling distortions, in some ways

perhaps akin to tinnitus in the case of phantomsia. There is certainly evidence of the need for more resources for these patients both in terms of direct therapeutic options and indirect supportive options.

Authorship in conjunction with the ICMJE criteria

Carl Philpott – 1, 2, 3, 4

Duncan Boak – 1, 2, 3

Joanne Dixon – 1, 3, 4

Conflict of Interest

None reported.

Funding

No funding was required for this study.

Reporting guidelines

This study has been reported in line with the Strobe guidelines

Acknowledgements

Thanks to Professor Thomas Hummel for inspiring the survey and providing input to the structure of the survey.

Tables

Table 1: Responses to question on parosmia impact (n/%)

Is your parosmia. . .	YES	NO	Not applicable
a daily occurrence?	65	9	26
intense?	44	27	29
having an impact on your enjoyment of every day	62	13	25
is causing you to (or has done) lose weight?	21	51	28

Table 2: Responses to question on phantomsia impact (n/%)

Is your phantomsia. . .	YES	NO	Not applicable
a daily occurrence?	31	31	38
intense?	33	25	42

Is your phantasmia. . .	YES	NO	Not applicable
having an impact on your enjoyment of every day	32	29	39
is causing you to (or has done) lose weight?	10	47	43

Table 3: Self-help measures tried by participants

Measure tried	Participants (%)
Valsalva (closing one’s mouth, pinching one’s nose shut while pressing out as if blowing up a balloon)	33
Head movements	20
Stimulating the nose with other smells	67
Stimulating the nose with water	39
Stimulating the nose with deep breaths in through the nose	53
Stimulating the nose with menthol/mustard/pepper spray	33

Table 4: Stimulants that provoke symptoms

Stimulant	Participants (%)
Smells	61
Flushing the nose	2
Pungent smells (trigeminal effect)	17
Sneezing	15
Changes in air pressure	25
Air travel	3
Physical exercise	5

References

1. Lechner M, Chandrasekharan D, Jumani K, et al. Anosmia as a presenting symptom of SARS-CoV-2 infection in healthcare workers – A systematic review of the literature, case series, and recommendations for clinical assessment and management. *Rhinology*. 2020;Accepted - in press.
2. Vaira LA, Salzano G, Deiana G, De Riu G. Anosmia and Ageusia: Common Findings in COVID-19 Patients. *Laryngoscope*. 2020.
3. Yan CH, Faraji F, Prajapati DP, Boone CE, DeConde AS. Association of chemosensory dysfunction and Covid-19 in patients presenting with influenza-like symptoms. *Int Forum Allergy Rhinol*. 2020.
4. Lechien JR, Chiesa-Estomba CM, De Siati DR, et al. Olfactory and gustatory dysfunctions as a clinical presentation of mild-to-moderate forms of the coronavirus disease (COVID-19): a multicenter European study. *Eur Arch Otorhinolaryngol*. 2020.
5. Moein ST, Hashemian SMR, Mansourafshar B, Khorram-Tousi A, Tabarsi P, Doty RL. Smell dysfunction: a biomarker for COVID-19. *Int Forum Allergy Rhinol*. 2020.
6. Parma V, Ohla K, Veldhuizen MG, et al. More than just smell - COVID-19 is associated with severe impairment of smell, taste, and chemesthesis. *medRxiv*. 2020:2020.2005.2004.20090902.

7. Menni C, Valdes AM, Freidin MB, et al. Real-time tracking of self-reported symptoms to predict potential COVID-19. *Nature Medicine*. 2020.
8. Hummel T, Whitcroft KL, Andrews P, et al. Position paper on olfactory dysfunction. *Rhinol Suppl*. 2017;54(26):1-30.
9. Damm M, Temmel A, Welge-Lüssen A, et al. [Olfactory dysfunctions. Epidemiology and therapy in Germany, Austria and Switzerland]. *HNO*. 2004;52(2):112-120.
10. Philpott CM. Smell and Taste Disorders in the UK: First experiences with a specialised smell and taste outpatient clinic. *Bulletin of the Royal College of Surgeons of England*. 2014;96:156-159.
11. Frasnelli J, Landis BN, Heilmann S, et al. Clinical presentation of qualitative olfactory dysfunction. *Eur Arch Otorhinolaryngol*. 2004;261(7):411-415.
12. Philpott CM, Boak D. The impact of olfactory disorders in the United kingdom. *Chem Senses*. 2014;39(8):711-718.
13. Erskine SE, Philpott CM. An unmet need: Patients with smell and taste disorders. *Clin Otolaryngol*. 2020;45(2):197-203.
14. Liu D, Sabha M, Damm M, et al. Parosmia is associated with relevant olfactory recovery after olfactory training. In. Authorea2020.
15. Reden J, Maroldt H, Fritz A, Zahnert T, Hummel T. A study on the prognostic significance of qualitative olfactory dysfunction. *Eur Arch Otorhinolaryngol*. 2007;264(2):139-144.
16. Saltagi MZ, Rabbani CC, Ting JY, Higgins TS. Management of long-lasting phantosmia: a systematic review. *Int Forum Allergy Rhinol*. 2018;8(7):790-796.
17. List of Abstracts from the Thirty-second Annual Meeting of the Association for Chemoreception Sciences. *Chemical Senses*. 2010;35(7):627-644.
18. Leopold D. Distortion of olfactory perception: diagnosis and treatment. *Chem Senses*. 2002;27(7):611-615.
19. Ball S, Boak D, Dixon J, Carrie S, Philpott C. Barriers to Effective Healthcare for Patients Who Have Smell or Taste Disorders. In. Medrxiv2020.
20. Pierre B, Paul A, Patrick F, David M. Distorted Odorant Perception: Analysis of a Series of 56 Patients With Parosmia. *Archives of Otolaryngology - Head & Neck Surgery*. 2005(2):107.