

Abdullah Al Noman¹, Md. Shofiqul Islam¹, Samiron Sana^{2,3}, Prapti Mondal⁴, Rima Islam Meem¹, Shohel Rana³, Debashish Mondol³, Manoshi Sana⁵, Sheikh I. Hossain⁶, Taufique Joarder⁷, and Kishor Mazumder^{3,8}

¹Department of Genetic Engineering & Biotechnology, Faculty of Biological Science and Technology, Jashore University of Science and Technology, Jashore 7408, Bangladesh

²Pharmacy Discipline, Life Science School, Khulna University, Khulna 9208, Bangladesh

³Department of Pharmacy, Faculty of Biological Science and Technology, Jashore University of Science and Technology, Jashore 7408, Bangladesh.

⁴Department of Zoology, Rajshahi College, Rajshahi 6100, Bangladesh.

⁵Dhaka Medical College, Dhaka 1000, Bangladesh.

⁶School of Mechanical and Mechatronic Engineering, University of Technology Sydney, 81 Broadway, Ultimo, NSW 2007, Australia.

⁷Executive Director, Public Health Foundation, Bangladesh

⁸School of Biomedical Science, Charles Sturt University, NSW 2678, Australia.

October 1, 2020

Abstract

Background: The ongoing acute respiratory disease pandemic termed COVID-19 caused by a newly emerged coronavirus has jeopardized the world's health and economic sectors. As of July 20, 2020, the virus now known as SARS-CoV-2 has already infected more than 14 million individuals and killed 612,815 patients with a mortality rate of 4.12% around the world regardless of age, gender and race.

Main body: Bangladesh has become one of its worst sufferers, with 207,453 infected people and 2,668 related deaths with a mortality rate of 1.29% until July 20, 2020. More than 50% of COVID-19 patients in Bangladesh are aged between 21 and 40 but elderly people aged more than 60 have the highest mortality rate (44.7%). Male individuals are also more susceptible to the virus than females and consist of 71% and 79% among the infected and deceased patients respectively. The most prevalent clinical features following the virus incubation period are fever, fatigue and dry cough. A phylogenetic analysis study elucidated that the virus strain found in the country has 9 single-nucleotide variants, mostly in the ORF1ab gene, and a sequence containing 3 successive variants in the N protein, which reflects a weaker strain of SARS-CoV-2, implicating a possibility of its lower mortality rate. Another investigation of 184 genome samples of SARS-CoV-2 across the country implicated a close homology with a European haplotype of SARS-CoV-2. The country has also joined the race of vaccine development and started phase-III clinical trials of a candidate vaccine developed by Sinovac Research and Development Co Ltd, China.

Conclusion: Bangladesh, as a developing country, still struggles with the pandemic and needs to scale up its response operation and improve health care facilities such as testing capacity, institutional quarantine and isolation centers and promote awareness. Preventive measures such as strict lockdown, social distancing and boosting the existing immune system are thus considered the only contrivances.

Background

In late December 2019, the world came to know about an unknown threat caused by a pathogen with unidentified etiology originating from a seafood market in Wuhan in Hubei Province, China, and the Chinese Center for Disease Control and Prevention (CCDC) proclaimed it as novel coronavirus pneumonia (NCP) [1]. The

virus now known as severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) is included in the orthocoronavirinae subfamily, which belongs to the Coronaviridae family and is genetically grouped into main genera: *Alphacoronavirus* (α -CoV), *Betacoronavirus* (β -CoV), *Deltacoronavirus* (δ -CoV) and *Gammacoronavirus* (γ -CoV) [2]. Seven species of coronavirus, including the latest being SARS-CoV-2, have been identified as pathogenic for humans [3]. To date, two severe disease outbreaks associated with severe acute respiratory syndrome coronavirus (SARS-CoV) and Middle East respiratory syndrome coronavirus (MERS-CoV) have occurred by coronaviruses in China (2002-03) and the Middle East (2012), respectively [4, 5]. On the other hand, four coronaviruses, namely, HCoV-229E, HCoV-NL63 of alpha-*genus* and HCoV-OC43, HCoV-HKU1 of beta *genus*, have been revealed to cause mild respiratory complications in humans [2]. The International Committee on Taxonomy of Viruses identified its symptomatic and biological similarity with SARS-CoV, hence renamed the virus the severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2), and the disease caused by this virus was renamed COVID-19 (coronavirus disease-19) by the World Health Organization (WHO). It is a positive stranded-RNA virus belonging to the sarbecovirus subgenus of the β -coronavirus, which also comprises other zoonotic viruses, such as SARS-CoV and bat SARS-like CoV [3]. Similar to other viruses, SARS-CoV-2 thrives on several intermediate and final hosts before infecting humans, which makes the prevention and treatment of viral infection more difficult. Although the virus has a lower mortality rate than SARS-CoV and MERS-CoV, it possesses high infectivity and transmissibility. A genome sequencing study of novel coronavirus showed that SARS-CoV-2 has 79.5% and 96% resemblance to SARS-CoV and SARSr-CoV-RaTG13 (bat SARS coronavirus), respectively, which implies the origin of coronavirus from a bat [6]. There have been several reports published on the discovery of a large number of SARS-related coronaviruses from bats, a natural reservoir of coronaviruses [7–9]. As of 20 July 2020, a total of 14,845,850 people have been infected, and 612,815 have died of the disease globally [10]. In Bangladesh, a total of 207,453 people have been infected by the virus and 2,668 have died of the disease as of 20 July 2020, with a mortality rate of 1.29%. This article aims to discuss the epidemiological study, genomic features, diagnosis, prevention and treatment scenario of Bangladesh.

Main Text

Composition of the Genome and Molecular Basis of Pathogenesis

The genome of SARS-CoV-2 varies from approximately 26 to 32 kilobases and contains 14 open reading frames (ORFs). The longest ORF at the 5' terminus that represents the major portion of the genome (67%) encodes 16 non-structural proteins, while the 3' terminus encodes structural proteins [11]. Of them, the membrane protein (M), nucleocapsid protein (N), envelope protein (E) and spike glycoprotein (S) are the four main structural proteins shown in **Figure 01** .




Figure 1: Structure of SARS-CoV-2.

Moreover, its genome comprises eight accessory proteins, namely, 3a, 3b, p6, 7a, 7b, 8b, 9b, and orf1 [12]. There is an ample amount of M glycoproteins in the SARS-CoV-2 genome that are responsible for the formation of virus particles and conduce virus assembly. It consists of a small $-NH_2$ terminal domain uncovered to the outside of the virion, three transmembrane domains and a much longer $-COOH$ terminus that resides inside the virion [13]. The N protein is the only protein present in the nucleocapsid. Although it is predominantly intricated in various viral genome-related processes, particularly in virus assembly and the formation of nucleocapsids, it is also involved in replication-transcription complexes in infected cells and host cellular responses to viral infection [14, 15]. The smallest structural protein E is involved in various processes related to virus replication cycles, such as virus assembly, budding, envelope formation and pathogenesis [16]. The S protein mainly paves the way for the virus to bind and enter into the host cell membrane and results in infection. It consists of two subunits: the S1 subunit, which contains a signal peptide and the receptor-binding domain, and the S2 subunit, which contains conserved fusion proteins, a transmembrane domain and the cytoplasmic domain [17]. Hoffmann M. et al. demonstrated that the S protein of the virus, similar to SARS-CoV, has a strong propensity to bind with human angiotensin-converting enzyme 2 (ACE2) and targets ACE2 as a receptor for entry and undergoes structural changes to merge with the host [18]. However, SARS-CoV-2 is more dangerous because its affinity towards ACE2 is more than 10-fold higher than that of SARS-CoV [19]. Furthermore, novel coronavirus also utilizes the cellular protease TMPRSS2 (transmembrane protease, serine 2) because the S protein needs to be primed first before entering host cells. Generally, SARS-CoV-2 has less sequence homology with SARS-CoV (approximately 79%) and MERS-CoV (approximately 50%) [20].

Analysis of the genome sequence conducted by Lu R. et al. showed that its genome possesses 88% sequence homology with two bat coronaviruses, bat-SL-CoVZC45 and bat-SL-CoVZXC21 [20]. All the genome sequencing data of SARS-CoV-2 from all over the world are being uploaded to the Global Initiative on Sharing All Influenza Data (GISAID), which facilitates the process of analyzing and understanding its genomic biology. A total of 71,000 genomic sequences were available on the website until 24 July 2020 [21]. Following the entry and denudation of the virus, its genome commences the transcription process followed by the translation process. The replication and transcription processes of coronaviruses occur in the cytoplasm. The mechanism of replication requires continuous RNA synthesis to determine whether transcription involves discontinuous synthesis [22]. These mechanisms are stimulated by the replication-transcription complex encoded by a 20 kb replicase gene and presumed to consist of as many as 16 viral proteins and different proteins

involved with cellular processes [23]. The expression of the replicase gene is stimulated by the translation of the genomic RNA. After an individual is exposed to a potential virus source such as bat or pangolins or persons infected by the virus, human antigen processing cells (APCs) and virus-specific T lymphocytes counter the entry of the virus. APC, along with human MHC (major histocompatibility complex) molecules known as human leukocyte antigens (HLAs), responds to and mediates defense against novel coronavirus attacks [24]. Genetic polymorphism in HLA genes elucidates why individual susceptibility to novel coronavirus varies in a diverse population, and this difference among individual susceptibility is also promoted by genetic polymorphisms in mannose-binding lectin (MBL).

The genome variants of SARS-COV-2 strain from Bangladesh

The first complete genome sequence of SARS-CoV-2 (CHRF_nCoV19_0001) from a Bangladeshi isolate was submitted to the GISAID database on 12 May 2020. Bangladesh based Child Health Research Foundation conducted the sequencing project [38]. As of July 17, genome sequencing of 222 samples from Bangladeshi SARS-CoV-2 isolates was submitted on the GISAID database, of which 173 samples were sequenced by the Bangladesh Council of Scientific and Industrial Research (BCSIR) [39]. Jashore University of Science and Technology had announced on 24 June 2020 that they had sequenced the complete genome of another three SARS-CoV-2 strain that is responsible for the respiratory infections in the southern part of the country [40]. A comparative study of the Bangladeshi strain and other strains around the world revealed that there were 9 single-nucleotide variants in the Bangladeshi strain, mostly in the ORF1ab gene and also a sequence containing 3 successive variants in the N protein which probably reflects a weaker strain of SARS-CoV-2 and explain the reason of low mortality rate in Bangladesh [41]. Phylogenetic analysis elucidated that there is sequence homology among the Bangladeshi strains and strains from Taiwan, Greece and Kazakhstan which implicates that the virus strains found in these countries were descendent from a weaker strain of the same origin, probably Michigan or Arizona in the United States [41]. Another investigation of 184 genome samples of SARS-CoV-2 across the country found 634 mutations located in the whole genome that results in 274 nonsynonymous substitutions in 22 different proteins. Among the spike protein variants circulating across the country, G614 is most prevalent followed by L323 (94%) in RNA- dependent RNA polymerase (RdRp), K203 (82%) and R204 (82%) in nucleocapsid, and F120 (78%) in NSP2. These mutations implicated a close homology with a European haplotype of SARS-CoV-2. These sequencing data will help to predict prognosis and develop an effective vaccine for the treatment of COVID-19 patients in Bangladesh.

Epidemiology of COVID-19 in Bangladesh

The novel coronavirus outbreak was announced as a public health emergency of international concern on the 30th of January, 2020 by the World Health Organization (WHO) and as a controllable pandemic on the 11th of March [17]. As of July 20, 2020, a total of 14,845,850 people had been infected by the virus and 612,815 people had died of the disease globally [10]. The worst-hit country being the USA with almost 4 million confirmed cases which constitutes almost 27% of the total cases and 143,834 fatalities. In Europe, Russia was the country with the highest number of confirmed cases by July 20, 2020; a total of 777,486 people were infected and 12,427 persons had died. The global distribution of confirmed COVID-19 cases has been illustrated in **Figure 02**. [10].




Figure 2: Global distribution of confirmed cases of COVID-19 pandemic

According to epidemiologic data, Yemen had the highest mortality rate around the world (27.7%) followed by France (17.3%) and UK (15.4%) until 20 July 2020 [25]. Since there is no identified preexisting immunity against the virus, the infection rate is increasing in an unpredictable manner and all individuals are vulnerable to the virus. One of the alarming news is that since SARS-CoV-2 is an RNA virus, it continuously mutates and even after taking preemptive precautions and measures, the number of infected cases is increasing.

Bangladesh has become the new hotspot of COVID-19 and has the highest number of infected patients among the South Asian countries after India and Pakistan. According to the Institute of Epidemiology, Disease Control and Research (IEDCR) so far 207,453 people have been infected and 2,668 have died of the disease in Bangladesh until 20 July 2020, with a mortality rate of 1.29%. Around the same time, 113,558 of COVID-19 patients have recovered from the disease with a recovery rate of 54.8% [26]. The geographical distribution of confirmed cases was available on 70% cases (144,281/207,453). Dhaka division has the highest number of infected patients in the country with more than 50% of all reported cases, of which 48,322 have been reported from the country’s capital Dhaka city. Chattogram (29,661) and Rajshahi (10,477) divisions positioned the second and third places regarding most confirmed COVID-19 patients, respectively [26]. Of the 2,668 deceased cases, the highest number was also reported from the Dhaka division (1,305), followed by the Chattogram division (673) and the Khulna division (173) (**Table 01**) [26, 27]. The country reported its first confirmed SARS-CoV-2 infection on 8 March and its first death on 18 March 2020. In a 24 hour period, the highest 4,019 cases were reported on 2 July and the highest 64 fatalities on June 30 [28]. It saw a rapid rise of infection at the middle of May; almost 90,038 confirmed cases and 1,136 deaths have been reported between week 21 to week 25 which is more than the total number combined in the initial days, albeit infection had been on a steady increase in the initial month [26]. The country reached 10,000 cases on the 3rd of May, exceeded 100,000 confirmed cases on 18 June and has recently crossed 200,000 infected patients on 18 July.

Table 01: COVID-19 Confirmed cases in Bangladesh (up to 20 July 2020)

Division	Total confirmed case	Percentage (%)	Total deaths	Percentage (%)
Dhaka	77,354	53.61	1,305	48.91
Chattogram	29,661	20.56	673	25.22

Division	Total confirmed case	Percentage (%)	Total deaths	Percentage (%)
Rajshahi	10,477	7.27	144	5.40
Khulna	9,483	6.58	173	6.49
Sylhet	6,672	4.62	125	4.69
Barisal	4,021	2.79	100	3.75
Mymensingh	3,687	2.55	58	2.17
Rangpur	2,926	2.02	90	3.37

n 20 July 2020, the COVID-19 attack rate i.e AR (the total number of cases/ the total population) in Bangladesh was 1218.1 per 1 million and all of the 64 districts have been infected by SARS-CoV-2. The highest AR was observed in the Dhaka division (3,182.6/1,000,000), the most populous division of the country and within the division the country capital Dhaka has the highest AR (almost 12,915.9/1,000,000) [29]. A study showed that the mean reproduction number of COVID-19 in Bangladesh until June 25, 2020, is 1.40 (95% CI 1.11, 1.73) [30]. Analysis of demographic data of COVID-19 patients elucidated that, though more than 50 % of patients infected by SARS-CoV-2 in Bangladesh are aged between 21 and 40; the highest mortality rate was observed in elderly people aged more than 60 years as they have a relatively weaker immune system (**Figure 03**) [26, 31].




Figure 3: Cases and deaths percentage by ages in Bangladesh.

Male individuals were more susceptible to the virus than females and consist of 71% and 29% confirmed cases, respectively (**Figure 04 A**). Among the deceased 2,104 were male and 564 were female [31]; the death ratio of males to females was 79:21 (**Figure 04 B**).




Figure 4: **A:** Cases percentage by gender **B:** Deaths percentage by gender

Out of the 116,226 cases with known outcomes in Bangladesh by 20 July 2020 mortality rate is almost 2.3% (2,668/116,226) which comparatively lower than the global averages 6.4% (612,815/9,514,219) [10, 26]. SARS-CoV-2 infection to the health professionals has been another problem for Bangladesh because there are only 93,358 MBBS doctors and 9,569 BDS doctors in the country according to the recent data of DGHS [32]. As of 18 July 2020, there were a total of 5,690 infected cases among healthcare professionals including 1,995 doctors, 1,536 nurses and 2,159 other personnel and of them, 67 doctors died from the disease [33]. Law enforcement personnel are also at risk of SARS-CoV-2 infection as they are engaged in maintaining government injunction and thus 13,316 policemen have been infected by the virus, and 53 of them have died till the 17th of July 2020. Yet 10,076 cops have already cured of the disease and returned to their respective station [34]. Rohingya refugees living in Cox's Bazar district are also vulnerable populations to infectious disease because more than 860,000 Rohingya refugees dwell in such a congested place [35]. As of 19 July 2020, a total of 62 COVID-19 confirmed cases and 6 related deaths have been reported in Rohingya camps. But the exact cases may be more because so far only 1,288 tests have been conducted among the population [36].

Clinical Features of COVID-19 among Bangladeshi patients

Though the incubation period of SARS-CoV-2 was reported to be 2-10 days according to WHO,[37] it demonstrated to infect people with a median incubation period of 3 days [38]. The maximum latency was observed up to 24 days which is longer compared to MERS (5 days) and seasonal flu (2 days). This implies the necessity for longer periods of quarantine or effective monitoring of persons potentially exposed to the pathogen [39]. The disease caused by the SARS-CoV-2 manifests with diverse clinical characteristics ranging from asymptomatic patients to acute pneumonia with multiorgan failure. After analyzing various symptoms, the disease is now classified into four levels of severity [40]. Mild patients only exhibit symptoms of viral infection in the upper respiratory tract while moderate patients manifested with fever, respiratory symptoms of cough and shortness of breath. Patients with severe cases manifest with dyspnea, acute respiratory distress

syndrome (ARDS) and septic shock. Finally, patients who need ICU support are considered as critical and show features such as respiratory arrest, septic shock, and multiple organ impairment [41].

A study of 103 RT-PCR confirmed COVID-19 Bangladeshi patients demonstrated that 74.76% were mild, 9.71% were moderate and 15.53% were severely ill patients. The most predominant clinical features include fever (78.6%), fatigue (68%) and cough, (44.7%) loss of appetite (37.9%), shortness of breath (37.9%) and anosmia (35.0%) [42]. Another study of 100 COVID-19 patients demonstrated symptoms like rhinitis (13.0%), body ache (13%), headache (12.0%), sputum (7.0%) abdominal pain (4.0%) hemoptysis (3.0%) [43]. A cohort study of 201 patients showed that very few patients were manifested with rare symptoms like burning body (1%), toothache (1%), itchiness (0.5%), red-eye (0.5%), oral ulcer (0.5%), constipation (0.5%) and 4.5% were asymptomatic [44]. Hypertension (34.0%) was the most prevalent comorbidities associated with the patients followed by diabetes mellitus (21.4%). Ischemic heart disease (9.7%), chronic kidney disease (7.8%) renal disease (8.0%), asthma/COPD (6.0%) were also manifested in the patients [42].

Prevention and Diagnosis scheme of COVID-19 in Bangladesh

The government of Bangladesh announced the first lockdown on March 26 and extended it to May 16 [45]. On March 16, the government announced to shut down all the educational institutions from March 17 to 31 and later declared to continue the closure up to September until the infection rate has considerably decreased [46]. After that, the government lifted the lockdown for a few days and saw a massive increase in both infected and death numbers. The government has planned to impose a zone-wise lockdown from the 9th of June under a pilot project. According to the plan, infected areas will be divided into 3 zones depending on COVID-19 severity; Red zones- strict lockdown will be maintained to decrease the spreading of the disease and no person is allowed to cross the zone, Yellow zones- limited restriction will be imposed on the regular life and Green Zones- where virtually no or very few COVID-19 patients exist [47]. The country suspended all international travel on April 5, except for flights to and from China [48]. The government could not impose strict social distancing measures, especially in Dhaka where lives almost 1.1 million slum dwellers who earn their day-to-day livelihood [49]. Moreover, the majority of these slum dwellers have no knowledge of the threat posed by COVID-19, and they live in an unhygienic condition with very little supply of water and one bathroom for every 10–16 families that made them more vulnerable to infectious disease [50].

Currently, 80 public and private laboratories (46 in Dhaka and 34 in outside of Dhaka) including several public universities that have Real-time PCR machines have been conducting COVID-19 tests [51, 52] Total of 1,041,661 tests were conducted by 20 July 2020, with an overall positivity rate of 20% and almost 61.3% (638,258/1,041,661) tests were done in the laboratories of the Dhaka city [28]. Although the number of testing has been elevating day by day, it is still too far from reaching the target number required to test every possible suspected case, and among the South-Asian countries, Bangladesh was placed only above Afghanistan regarding COVID-19 testing. Only 6,322 per 1 million testings have been done until 20 July 2020 (**Figure 05**) [10, 53].




Figure 5: Test per 1 million people in the South Asian countries.

Another diagnostic testing kit was developed by Gonoshasthaya Kendra, Bangladesh on 17 March 2020. This cost-effective kit (only \$3.25 to produce) named ‘Gonoshasthaya Rapid Dot Blot’ can test antigen as well as antibody using saliva and swab samples and yields results within 15 minutes. The team led by Dr. Bijon Kumar Sil had been waiting for approval from the Directorate General of Drug Administration (DGDA) to use their kit immediately [54, 55]. In recent clinical trials, the Gonoshasthaya’s kit has shown 97.7% sensitivity and 96% specificity in internal validation that satisfies the US FDA guidelines for antibody testing kits [56].

Treatments scenario of COVID-19 in Bangladesh

So far, 91,229 patients are currently under treatment in different medical institutes, and 43,007 patients were placed in isolation units as of July 20, 2020. Currently, the country has 629 institutional quarantine centers that can accommodate 31,991 patients across 64 districts, and as of 20 July 23,331 individuals were gone under institutional quarantine and of them, 17,406 have got release while 335,643 out of 389,150 individuals have been released from home quarantine [53]. Different antiviral drugs and treatment strategies have been implemented to treat COVID-19 patients. DGHS recommended controversial drugs such as chloroquine and hydroxychloroquine for the treatment in the “National Guidelines on Clinical Management of Coronavirus Disease-2019” [57]. A medical team from Bangladesh Medical College Hospital claimed that a combination of antiprotozoal medicine Ivermectin and antibiotic Doxycycline was effective for the treatment of 60 COVID-19 patients [58]. Clinical trials are going on and until now 400-500 COVID-19 patients have received the drugs, of them almost 98% of patients had recovered within 4-14 days [59]. Bangladesh based Beximco Pharmaceuticals had launched the world’s first generic Remdesivir-an antiviral drug for the treatment of COVID-19 patients on 21 May 2020 [60]. In a recent clinical trial, it has demonstrated potential effectivity against SARS-CoV-2 and lessened the recovery time [61]. Remdesivir which works by inhibiting the replication of SARS-CoV-2 has now become the talk of the town and recently received “Emergency Use Authorisation” from the FDA and developed by Gilead Sciences [62]. Bangladeshi based

pharmaceutical Globe Biotech Limited has announced on 1st July 2020 that they are developing a COVID-19 vaccine which showed effectivity on preliminary trials on animal models [63]. They have also developed a guideline for clinical trials which will be conducted after getting approval from the Bangladesh Medical Research Council (BMRC) and claimed to produce the vaccine within six months after clinical trials. The research group of 12 scientists analyzed 5743 genome sequences of the SARS-CoV-2 from the US National Center for Biotechnology Information (NCBI) through the bioinformatics approach and conducted the test through the Surface plasmon resonance (SPR) method to detect molecular interactions [64, 65]. The BMRC has recently approved the phase-III clinical trials of a candidate vaccine developed by Sinovac Research and Development Co Ltd, China in Bangladesh. The trials will involve 4200 volunteers at seven different hospitals and will begin in the early of August this year under the supervision of the International Centre for Diarrhoeal Disease Research, Bangladesh (icDDR,b) [66, 67]. The country has recently started clinical trials of convalescent plasma therapy (CPT) and received ethical approval from DGDA and the BMRC. 18 patients were administered plasma from recovered patients and the government in collaboration with several national organization has recently inaugurated a plasma network namely ‘Shohojodha’ to consolidate plasma therapy across the country [68, 69].

Economic impact of COVID-19 on Bangladesh

Bangladesh likewise the most countries of the world are facing the gravest economic crisis for this unprecedented pandemic. Export Promotion Bureau of Bangladesh (EPB) revealed that export earnings have fallen to 520 million USD in March this year from 3.03 billion USD in the same month of the previous year [70]. The International Monetary Fund (IMF) had projected the country’s GDP growth to be 2%-3.8% for the fiscal year 2019-2020 from 7.4%-8.2% which was estimated before [71]. According to the Bangladesh Bureau of Statistics (BBS), currently thirty-four million people live below the poverty line that constitutes 20.5% of the total population. A survey conducted by South Asian Network on Economic Modelling (SANEM) denoted that if the income level for the poverty line is increased 1.25 percent the number of poor people would be raised to thirty-six million [72]. The prime minister of Bangladesh had declared to provide 19 incentive packages of Tk 1.03 trillion (equivalent to 12.13 billion USD) to cushion the economic blow of the COVID-19 pandemic [73]. She also announced an incentive package of 600 million dollars to facilitate the industries dealing with exporting products [74]. But the government alone cannot confront this unprecedented situation and requires both public and private collaboration. Co-operation between the local as well as international organizations such as the World Health Organization and the World Economic Forum is mandatory to combat this massive economic blow caused by COVID-19.

Points to be considered in Bangladesh

- Zone-wise lockdown should be continued, and the government should keep in mind both the economic challenges and COVID-19 infection rate.
- Implementation of a nationwide contact tracing is mandatory, which will help to locate the infected individuals, their contacted persons and prevent secondary infections.
- Both the national and international public health agencies should come together for scaling up surveillance operations throughout the country.
- Uses of low-cost and rapid diagnostic kits (such as Gonoshasthay’s kit) would be an effective strategy to identify all possible suspected cases which will mitigate the spreading.
- Finally, raising campaigns about using face masks, maintaining social isolation and eliminating the rumors of COVID-19 among the mass population can be some possible measures.

Conclusion

The ongoing COVID-19 pandemic caused by this newly emerged virus is undoubtedly a matter of international health concern. Investigation of viral genomic, pathogenesis and epidemiological study have been continued to find potential treatment strategies. In Bangladesh, delay of strategic planning, scarcity of

health care facilities (PPE, ventilators), shortage of appropriate preventive measures (quarantine and isolation unit), emergency service, unawareness and lack of knowledge of the people about the COVID-19 is aggravating the situation. As a developing country, Bangladesh should focus on scaling up its readiness and response operation as well as improving the health care facilities such as upgrading the capacity of the testing laboratories, institutional quarantine centers and stocking personal protective equipment. Promoting knowledge and awareness of the COVID-19 among the mass people is also significant.

LIST of ABBREVIATIONS

COVID-19: Coronavirus Disease 2019; SARS-CoV-2: Severe Acute Respiratory Syndrome Coronavirus 2; HCoV-229E: Human Coronavirus 229E; HCoV-OC43: Human Coronavirus OC43; HCoV-NL63: Human Coronavirus NL63; HCoV-HKU1: Human Coronavirus HKU1; SARS-CoV: Severe Acute Respiratory Syndrome Coronavirus; MERS-CoV: Middle East Respiratory Syndrome Coronavirus; WHO: World Health Organization; SARS: Severe Acute Respiratory Syndrome; MERS: Middle East Respiratory Syndrome; ACE2: Angiotensin-Converting Enzyme 2; TMPRSS2: Transmembrane Protease, Serine 2; GSAID: Global Initiative on Sharing All Influenza Data; MHC: Major Histocompatibility Complex; SNP: Single Nucleotide Polymorphism; ARDS: Acute Respiratory Distress Syndrome; CT: Computed Tomography; RT-PCR: Reverse-Transcription Polymerase Chain Reaction; CDC: Centers for Disease Control and Prevention; IEDCR: Institute of Epidemiology, Disease Control and Research; DGHS: Directorate General of Health Services; AR: Attack Rate; DGDA: Directorate General of Drug Administration; BMRC: Bangladesh Medical Research Council; EBP: Extracorporeal blood purification.

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and material

All data and materials are available upon request

Competing interests

The authors declare that they have no competing interests

Funding

This work is managed to perform with the individual funding of all authors.

Authors' Contributions

All authors (AAN, MSI, SS, PM, RIM, SR, DM, MS, SIH, TJ, KM) contributed to the manuscript. AAN, MSI, SS organized the project. Validation investigation, resources, data curation and writing were done by all authors. SIH, TJ, KM contributed to the review. All authors read and approved the manuscript.

Acknowledgment

The authors are grateful to Dr. Niranjana Kumar Sana, Professor, Department of Biochemistry and Molecular Biology, University of Rajshahi, Rajshahi 6205, Bangladesh and Dr. Asish Kumar Das, Professor, Pharmacy Discipline, Khulna University, Khulna 9208, Bangladesh for providing their thoughts and guidance.

REFERENCES

1. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020;395:497–506. doi:10.1016/S0140-6736(20)30183-5.

2. Chan JFW, To KKW, Tse H, Jin DY, Yuen KY. Interspecies transmission and emergence of novel viruses: Lessons from bats and birds. *Trends in Microbiology*. 2013;21:544–55. doi:10.1016/j.tim.2013.05.005.
3. Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med*. 2020;382:727–33. doi:10.1056/NEJMoa2001017.
4. Zhong NS, Zheng BJ, Li YM, Poon LLM, Xie ZH, Chan KH, et al. Epidemiology and cause of severe acute respiratory syndrome (SARS) in Guangdong, People’s Republic of China, in February, 2003. *Lancet*. 2003;362:1353–8. doi:10.1016/S0140-6736(03)14630-2.
5. Zaki AM, van Boheemen S, Bestebroer TM, Osterhaus ADME, Fouchier RAM. Isolation of a Novel Coronavirus from a Man with Pneumonia in Saudi Arabia. *N Engl J Med*. 2012;367:1814–20. doi:10.1056/NEJMoa1211721.
6. Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet*. 2020;395:507–13. doi:10.1016/S0140-6736(20)30211-7.
7. Li W, Shi Z, Yu M, Ren W, Smith C, Epstein JH, et al. Bats Are Natural Reservoirs of SARS-Like Coronaviruses. *Science* (80-). 2005;310.
8. Ge XY, Li JL, Yang X Lou, Chmura AA, Zhu G, Epstein JH, et al. Isolation and characterization of a bat SARS-like coronavirus that uses the ACE2 receptor. *Nature*. 2013;503:535–8. doi:10.1038/nature12711.
9. Yang L, Wu Z, Ren X, Yang F, He G, Zhang J, et al. Novel SARS-like betacoronaviruses in bats, China, 2011. *Emerg Infect Dis*. 2013;19:989–91. doi:10.3201/eid1906.121648.
10. Worldometer. Coronavirus Cases. Worldometer. 2020;:1–22. doi:10.1101/2020.01.23.20018549V2.
11. Khailany RA, Safdar M, Ozaslan M. Genomic characterization of a novel SARS-CoV-2. *Gene Reports*. 2020;19:100682. doi:10.1016/j.genrep.2020.100682.
12. Wu A, Peng Y, Huang B, Ding X, Wang X, Niu P, et al. Genome Composition and Divergence of the Novel Coronavirus (2019-nCoV) Originating in China. *Cell Host Microbe*. 2020;27:325–8. doi:10.1016/j.chom.2020.02.001.
13. Hogue BG, Machamer CE. Coronavirus Structural Proteins and Virus Assembly. In: *Nidoviruses*. American Society of Microbiology; 2014. p. 179–200. doi:10.1128/9781555815790.ch12.
14. de Haan CAM, Kuo L, Masters PS, Vennema H, Rottier PJM. Coronavirus Particle Assembly: Primary Structure Requirements of the Membrane Protein. *J Virol*. 1998;72:6838–50. doi:10.1128/jvi.72.8.6838-6850.1998.
15. Verheije MH, Hagemeijer MC, Ulasli M, Reggiori F, Rottier PJM, Masters PS, et al. The Coronavirus Nucleocapsid Protein Is Dynamically Associated with the Replication-Transcription Complexes. *J Virol*. 2010;84:11575–9. doi:10.1128/jvi.00569-10.
16. Schoeman D, Fielding BC. Coronavirus envelope protein: Current knowledge. *Virology Journal*. 2019;16. doi:10.1186/s12985-019-1182-0.
17. Abduljalil JM, Abduljalil BM. Epidemiology, genome, and clinical features of the pandemic SARS-CoV-2: a recent view. *New Microbes and New Infections*. 2020;35:100672.
18. Hoffmann M, Kleine-Weber H, Krüger N, Müller M, Drosten C, Pöhlmann S. The novel coronavirus 2019 (2019-nCoV) uses the SARS-coronavirus receptor ACE2 and the cellular protease TMPRSS2 for entry into target cells. *bioRxiv*. 2020.
19. Yuan Y, Cao D, Zhang Y, Ma J, Qi J, Wang Q, et al. Cryo-EM structures of MERS-CoV and SARS-CoV spike glycoproteins reveal the dynamic receptor binding domains. *Nat Commun*. 2017;8:15092.

20. Lu R, Zhao X, Li J, Niu P, Yang B, Wu H, et al. Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. *Lancet*. 2020;395:565–74.
21. GISAID - Initiative. <https://www.gisaid.org/>. Accessed 24 Jul 2020.
22. Sola I, Almazán F, Zúñiga S, Enjuanes L. Continuous and Discontinuous RNA Synthesis in Coronaviruses. *Annu Rev Virol*. 2015;2:265–88. doi:10.1146/annurev-virology-100114-055218.
23. Mousavizadeh L, Ghasemi S. Genotype and phenotype of COVID-19: Their roles in pathogenesis. *Journal of Microbiology, Immunology and Infection*. 2020. doi:10.1016/j.jmii.2020.03.022.
24. COVID-19 and SARS-CoV-2: Molecular Genetics Perspectives. https://www.researchgate.net/publication/340685184_COVID-19_and_SARS-CoV-2_Molecular_Genetics_-_Perspectives. Accessed 13 Jul 2020.
25. Mortality Risk of COVID-19 - Statistics and Research - Our World in Data. <https://ourworldindata.org/mortality-risk-covid#the-case-fatality-rate>. Accessed 21 Jul 2020.
26. IEDCR. <https://www.iedcr.gov.bd/>. Accessed 21 Jul 2020.
27. Bangladesh's Covid-19 death toll stands at 2,668. <https://unb.com.bd/category/Bangladesh/bangladeshs-covid-19-death-toll-stands-at-2688/54905>. Accessed 21 Jul 2020.
28. Coronavirus Disease 2019 (COVID-19) Information Bangladesh | corona.gov.bd. <https://corona.gov.bd/>. Accessed 21 Jul 2020.
29. Coronavirus disease (COVID-2019) Bangladesh situation reports-21. [https://www.who.int/bangladesh/emergencies/coronavirus-disease-\(covid-19\)-update/coronavirus-disease-\(covid-2019\)-bangladesh-situation-reports](https://www.who.int/bangladesh/emergencies/coronavirus-disease-(covid-19)-update/coronavirus-disease-(covid-2019)-bangladesh-situation-reports). Accessed 21 Jul 2020.
30. Estimation of Reproduction Number of COVID-19 in case of Bangladesh. https://www.researchgate.net/publication/342260886_Estimation_of_Reproduction_Number_of_COVID-19_in_case_of_Bangladesh. Accessed 20 Jul 2020.
31. Bangladesh crosses Germany in Covid-19 infections | Dhaka Tribune. <https://www.dhakatribune.com/bangladesh/2020/07/20/covid-19-50-die-2-928-new-cases-in-24hrs>. Accessed 21 Jul 2020.
32. COVID-19 in Bangladesh - Doctors, nurses getting sick: Protective gear, curbs on non-essential services can save lives | The Business Standard. <https://tbsnews.net/coronavirus-chronicle/covid-19-bangladesh/doctors-nurses-getting-sick-protective-gear-curbs-non>. Accessed 13 Jun 2020.
33. Two more doctors die of COVID-19 in Bangladesh. <https://www.newagebd.net/article/111476/two-more-doctors-die-of-covid-19-in-bangladesh>. Accessed 19 Jul 2020.
34. Coronavirus: 10,076 recover out of 13,316 infected policemen; 53 dead. <http://unb.com.bd/category/Bangladesh/coronavirus-10076-recover-out-of-13316-infected-policemen-53-dead/54785>. Accessed 19 Jul 2020.
35. UNHCR Bangladesh - Operational Dashboard : 2020 Indicators Monitoring (As of May 2020). <https://data2.unhcr.org/en/documents/details/77873>. Accessed 22 Jul 2020.
36. Emergency: Rohingya crisis - WHO Bangladesh situation report #15, 22 July 2020. <https://reliefweb.int/report/bangladesh/emergency-rohingya-crisis-who-bangladesh-situation-report-15-22-july-2020-period>. Accessed 25 Jul 2020.
37. Coronavirus Disease (COVID-19) Situation Reports-7. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports>. Accessed 27 Jan 2020.

38. Guan W, Ni Z, Hu Y, Liang W, Ou C, He J, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med.* 2020;382:1708–20. doi:10.1056/NEJMoa2002032.
39. Coronavirus Incubation Period (COVID-19) - Worldometer. <https://www.worldometers.info/coronavirus/coronavirus-incubation-period/>. Accessed 14 Jun 2020.
40. Wang Y, Wang Y, Chen Y, Qin Q. Unique epidemiological and clinical features of the emerging 2019 novel coronavirus pneumonia (COVID-19) implicate special control measures. *Journal of Medical Virology.* 2020;92:568–76. doi:10.1002/jmv.25748.
41. Cascella M, Rajnik M, Cuomo A, Dulebohn SC, Di Napoli R. Features, Evaluation and Treatment Coronavirus (COVID-19). StatPearls Publishing; 2020. <http://www.ncbi.nlm.nih.gov/pubmed/32150360>. Accessed 14 Jul 2020.
42. Shahed Morshed M, Al Mosabbir A, Chowdhury P, Mohammad S, Sorowar Hossain M. Clinical manifestations of patients with Coronavirus Disease 2019 (COVID- 19) attending at hospitals in Bangladesh. medRxiv. 2020;:2020.07.30.20165100. doi:10.1101/2020.07.30.20165100.
43. Mowla SGM, Azad KAK, Kabir A, Biswas S, Islam MR, Banik GC, et al. Clinical Profile of 100 Confirmed COVID-19 Patients Admitted in Dhaka Medical College Hospital, Dhaka, Bangladesh. *J Bangladesh Coll Physicians Surg.* 2020;:29–36. doi:10.3329/jbcp.v38i0.47445.
44. Ahmed NU, Islam MA, Kabir MA, Rahman MH, Sadat SA. Clinico-Pathological Findings of Bangladeshi Covid 19 Patients with their Clinical Outcome: Study of A Cohort of 201 Cases. *J Bangladesh Coll Physicians Surg.* 2020;:37–42. doi:10.3329/jbcp.v38i0.47346.
45. Coronavirus: Is Bangladesh putting its economy before people? | Humanitarian crises | Al Jazeera. <https://www.aljazeera.com/programmes/thestream/2020/04/coronavirus-bangladesh-putting-economy-people-200430180547872.html>. Accessed 5 Jun 2020.
46. Coronavirus Update in Bangladesh: Educational institutions may remain closed till September if virus does not fade: PM | The Business Standard. <https://tbsnews.net/bangladesh/education/all-educational-institutions-remain-closed-till-september-pm-74221>. Accessed 12 Jun 2020.
47. Bangladesh set to test zone-wise lockdown from Tuesday night | Dhaka Tribune. <https://www.dhakatribune.com/health/coronavirus/2020/06/09/bangladesh-set-to-test-zone-wise-lockdown-from-tuesday-night>. Accessed 12 Jun 2020.
48. Bangladesh: Government extends existing COVID-19 restrictions until April 25 and imposes curfew from April 10 /update 7. <https://www.garda.com/crisis24/news-alerts/331331/bangladesh-government-extends-existing-covid-19-restrictions-until-april-25-and-imposes-curfew-from-april-10-update-7>. Accessed 10 May 2020.
49. Bangladesh Bureau of Statistics. Slum Census and Floating population 2014. Dhaka (2015). <http://203.112.218.65:8008/PageWebMenuContent.aspx?MenuKey=423>. Accessed 18 Jun 2020.
50. Arias-Granada Y, Haque SS, Joseph G, Yanez-Pagans M. Water and Sanitation in Dhaka Slums: Access, Quality, and Informality in Service Provision. The World Bank; 2018. doi:10.1596/1813-9450-8552.
51. Why did DU discontinue Covid-19 testing? | Dhaka Tribune. <https://www.dhakatribune.com/health/coronavirus/2020/06/05/du-discontinues-covid-19-testing-amid-worsening-conditions>. Accessed 9 Jun 2020.
52. Directorate General of Health Services (DGHS). <https://www.dghs.gov.bd/index.php/en/component/content/article?id=> Accessed 21 Jul 2020.
53. COVID-19. <http://103.247.238.92/webportal/pages/covid19.php>. Accessed 21 Jul 2020.

54. Are Bangladesh's low-cost coronavirus test kits reliable? | Asia | An in-depth look at news from across the continent | DW | 17.04.2020. <https://www.dw.com/en/are-bangladeshs-low-cost-coronavirus-test-kits-reliable/a-53163373>. Accessed 1 Jun 2020.

55. Gonoshasthaya Kendra's rapid test kit performance trial report in limbo. <https://www.newagebd.net/article/106845/gonoshasthaya-kendras-rapid-test-kit-performance-trial-report-in-limbo>. Accessed 1 Jun 2020.

56. Covid-19: Gonoshasthaya Kendra takes second shot at external validation of antibody test kit | Dhaka Tribune. <https://www.dhakatribune.com/bangladesh/government-affairs/2020/07/05/gonoshasthaya-covid-19-test-kit-97-7-accurate-to-undergo-external-validation>. Accessed 7 Jul 2020.

57. National Guidelines on Clinical Management of Coronavirus Disease-2019". https://www.google.com/search?sxsrf=ALeKk02q_jUU5nbrnz4jei-d7t9zi-fVoA%3A1594800604375&ei=3LkOX5qlFsXt9QOzi5Fg&q=National+Guidelines+on+Clinical+Management+of+Coronavirus+Disease-2019.&oq=National+Guidelines+on+Clinical+Management+of+Coronavirus+Disease-. Accessed 6 Jun 2020.

58. Bangladeshi doctors claim to have found effective drugs to treat COVID-19 | The Business Standard. <https://tbsnews.net/coronavirus-chronicle/covid-19-bangladesh/bangladeshi-doctors-claim-have-found-effective-drugs-treat>. Accessed 3 Jun 2020.

59. Dr Tarek Alam: Unleashing the potential of Ivermectin. <https://unb.com.bd/category/special/dr-tarek-alam-unleashing-the-potential-of-ivermectin/54846>. Accessed 20 Jul 2020.

60. Beximco Pharma introduces world's first generic Remdesivir for COVID-19 treatment. <https://thefinancialexpress.com.bd/health/beximco-pharma-introduces-worlds-first-generic-remdesivir-for-covid-19-treatment-1590062827>. Accessed 2 Jun 2020.

61. Beigel JH, Tomashek KM, Dodd LE, Mehta AK, Zingman BS, Kalil AC, et al. Remdesivir for the Treatment of Covid-19 — Preliminary Report. *N Engl J Med*. 2020. doi:10.1056/nejmoa2007764.

62. Ledford H. Hopes rise for coronavirus drug remdesivir. *Nature*. 2020.

63. Bangladeshi firm Globe Biotech in race to develop Covid-19 vaccine. <https://thefinancialexpress.com.bd/trade/bangladeshi-firm-globe-biotech-in-race-to-develop-covid-19-vaccine-1593750347>. Accessed 4 Jul 2020.

64. Bangladeshi Globe Biotech also joins global race to develop Covid-19 vaccine. <https://en.somoynews.tv/9364/news/Bangladeshi-Globe-Biotech-also-joins-global-race-to-develop-Covid-19-vaccine>. Accessed 4 Jul 2020.

65. Globe Biotech claims to be the first Bangladeshi company to develop Covid-19 vaccine | Dhaka Tribune. <https://www.dhakatribune.com/bangladesh/2020/07/02/globe-biotech-to-become-the-first-bangladeshi-company-to-develop-covid-19-vaccine>. Accessed 4 Jul 2020.

66. Bangladesh to host late-stage trial of China's Sinovac COVID-19 vaccine - Reuters. <https://www.reuters.com/article/us-health-coronavirus-bangladesh/bangladesh-allows-late-stage-trial-of-chinas-sinovac-covid-19-vaccine-idUSKCN24L0KO>. Accessed 20 Jul 2020.

67. BMRC approves icddr,b in human trial of Chinese Covid-19 vaccine in Bangladesh. <https://www.thedailystar.net/bmrc-approves-icddrb-human-trial-chinese-covid-19-vaccine-in-bangladesh-1932893>. Accessed 20 Jul 2020.

68. Bangladesh experts advise against using unproven drugs to treat COVID-19 - bdnews24.com. <https://bdnews24.com/health/2020/06/01/bangladesh-experts-advise-against-using-unproven-drugs-to-treat-covid-19>. Accessed 3 Jun 2020.

69. ICT-driven PPP to fight Covid-19. <https://thefinancialexpress.com.bd/economy/bangladesh/ict-driven-ppp-to-fight-covid-19-1594909956>. Accessed 20 Jul 2020.
70. Coronavirus: Economy down, poverty up in Bangladesh | Asia | An in-depth look at news from across the continent | DW | 10.06.2020. <https://www.dw.com/en/coronavirus-economy-down-poverty-up-in-bangladesh/a-53759686>. Accessed 12 Jun 2020.
71. COVID-19 Impact on Bangladesh Economy. <https://www.arx.cfa/en/research/2020/06/soc290620-covid-19-impact-on-bangladesh-economy>. Accessed 24 Jun 2020.
72. Economic ramifications of Covid-19 in Bangladesh. <https://thefinancialexpress.com.bd/views/columns/economic-ramifications-of-covid-19-in-bangladesh-1588779315>. Accessed 13 Jun 2020.
73. Hopes still abound for economic turnaround. <https://thefinancialexpress.com.bd/views/columns/hopes-still-abound-for-economic-turnaround-1591458234>. Accessed 12 Jun 2020.
74. Covid-19: Economic challenges facing Bangladesh. <https://thefinancialexpress.com.bd/views/covid-19-economic-challenges-facing-bangladesh-1592064588>. Accessed 13 Jun 2020.